

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau**B4**

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/435, 14/00, 7/06, C12N 5/10, 15/10, 15/11, 15/12, 15/63, A61K 38/16</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/09552</b> (43) International Publication Date: <b>24 February 2000 (24.02.00)</b>																											
(21) International Application Number: <b>PCT/US99/18298</b> (22) International Filing Date: <b>13 August 1999 (13.08.99)</b>  (30) Priority Data: <table><tr><td>60/096,622</td><td>14 August 1998 (14.08.98)</td><td>US</td></tr><tr><td>60/096,815</td><td>17 August 1998 (17.08.98)</td><td>US</td></tr><tr><td>60/099,229</td><td>4 September 1998 (04.09.98)</td><td>US</td></tr><tr><td>60/105,368</td><td>23 October 1998 (23.10.98)</td><td>US</td></tr><tr><td>60/115,234</td><td>8 January 1999 (08.01.99)</td><td>US</td></tr><tr><td>60/119,931</td><td>12 February 1999 (12.02.99)</td><td>US</td></tr><tr><td>60/120,575</td><td>18 February 1999 (18.02.99)</td><td>US</td></tr><tr><td>60/132,020</td><td>30 April 1999 (30.04.99)</td><td>US</td></tr><tr><td>Not furnished</td><td>11 August 1999 (11.08.99)</td><td>US</td></tr></table> (71) Applicant: <b>GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US).</b>		60/096,622	14 August 1998 (14.08.98)	US	60/096,815	17 August 1998 (17.08.98)	US	60/099,229	4 September 1998 (04.09.98)	US	60/105,368	23 October 1998 (23.10.98)	US	60/115,234	8 January 1999 (08.01.99)	US	60/119,931	12 February 1999 (12.02.99)	US	60/120,575	18 February 1999 (18.02.99)	US	60/132,020	30 April 1999 (30.04.99)	US	Not furnished	11 August 1999 (11.08.99)	US	(72) Inventors: <b>JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). McCOY, John, M.; 56 Howard Street, Reading, MA 01867 (US). LaVALLIE, Edward, R.; 133 Ann Lee Road, Harvard, MA 01451 (US). COLLINS-RACIE, Lisa, A.; 124 School Street, Acton, MA 01720 (US). EVANS, Cheryl; 18801 Bent Willow Circle, Germantown, MA 20874 (US). MERBERG, David; 2 Orchard Drive, Acton, MA 01720 (US). TREACY, Maurice; 12 Foxrock Court, Dublin 4 (IE). AGOSTINO, Michael, J.; 26 Wolcott Avenue, Andover, MA 01810 (US). STEININGER, Robert, J., II; 100 Reed Street, Cambridge, MA 02140 (US). SPAULDING, Vikki; 11 Meadowbank Road, Billerica, MA 01821 (US). WONG, Gordon, G.; 239 Clark Road, Brookline, MA 02146 (US). CLARK, Hilary, F.; 495 Harkness Avenue, San Francisco, CA 94134 (US). FECHTEL, Kim; 46 Marion Road, Arlington, MA 02174 (US).</b>  (74) Agent: <b>SPRUNGER, Suzanne, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).</b>  (81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report.</i>	
60/096,622	14 August 1998 (14.08.98)	US																												
60/096,815	17 August 1998 (17.08.98)	US																												
60/099,229	4 September 1998 (04.09.98)	US																												
60/105,368	23 October 1998 (23.10.98)	US																												
60/115,234	8 January 1999 (08.01.99)	US																												
60/119,931	12 February 1999 (12.02.99)	US																												
60/120,575	18 February 1999 (18.02.99)	US																												
60/132,020	30 April 1999 (30.04.99)	US																												
Not furnished	11 August 1999 (11.08.99)	US																												
(54) Title. <b>SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM</b>																														
(57) Abstract  <b>Novel polynucleotides and the proteins encoded thereby are disclosed.</b>																														

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



5

## SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

10

This application is a continuation-in-part of the following applications:

- (1) provisional application Ser. No. 60/096,622 (GI 6075), filed August 14, 1998;
  - (2) provisional application Ser. No. 60/096,815 (GI 6076), filed August 17, 1998;
  - (3) provisional application Ser. No. 60/099,229 (GI 6077), filed September 4, 1998;
  - 15 (4) provisional application Ser. No. 60/105,368 (GI 6078), filed October 23, 1998;
  - (5) provisional application Ser. No. 60/115,234 (GI 6079), filed January 8, 1999;
  - (6) provisional application Ser. No. 60/119,931 (GI 6080), filed February 12, 1999;
  - (7) provisional application Ser. No. 60/120,575 (GI 6081), filed February 18, 1999;
  - (8) provisional application Ser. No. 60/132,020 (GI 6082), filed April 30, 1999;
  - 20 (9) provisional application Ser. No. 60/XXX,XXX (GI 6083), filed August 11, 1999;
- all of which are incorporated by reference herein.

25

FIELD OF THE INVENTION

The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins.

30

35

### BACKGROUND OF THE INVENTION

Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins and the polynucleotides encoding them that the present invention is directed.

### SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 120 to nucleotide 821;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 1 to nucleotide 1625;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;

- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825;
- 5 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
- 10 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 15 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:1.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821; the nucleotide sequence of SEQ ID NO:1 from nucleotide 120 to nucleotide 821; the nucleotide sequence of SEQ ID NO:1 from nucleotide 1 to nucleotide 1625; the nucleotide sequence of the full-length protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825; or the
- 25 nucleotide sequence of a mature protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825. In further preferred
- 30 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having

biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:2.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:1.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:1, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:1; and

(ab) the nucleotide sequence of the cDNA insert of clone  
co62\_12 deposited under accession number ATCC 98825;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

(bb) the nucleotide sequence of the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:1 to

a nucleotide sequence corresponding to the 3' end of SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821, and extending  
5 contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
10 NO:1 from nucleotide 120 to nucleotide 821, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 120 to nucleotide 821, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 120 to nucleotide 821. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 1 to nucleotide 1625, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 1 to nucleotide 1625, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 1 to nucleotide 1625.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) a fragment of the amino acid sequence of SEQ ID NO:2, the  
25 fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
- (c) the amino acid sequence encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:2. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:3 from nucleotide 96 to nucleotide 1013;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the  
15 cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
20 insert of clone lo311\_8 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:3.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013; the nucleotide sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013; the nucleotide sequence of the full-length protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825; or the  
5 nucleotide sequence of a mature protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825. In further preferred  
10 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from amino acid 162  
15 to amino acid 171 of SEQ ID NO:4.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:3.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (aa) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and
    - (ab) the nucleotide sequence of the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bb) the nucleotide sequence of the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

10 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:3 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and



(c) the amino acid sequence encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from  
10 amino acid 162 to amino acid 171 of SEQ ID NO:4.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825;
- 25 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:5.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825; the nucleotide sequence of the full-length  
10 protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825. In further preferred  
15 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having  
20 biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:6.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:5.

Further embodiments of the invention provide isolated polynucleotides produced  
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(ab) the nucleotide sequence of the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(bb) the nucleotide sequence of the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5, and extending  
20 contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:5 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825, and extending  
25 contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ  
10 ID NO:6 having biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pj193\_5 deposited under accession  
20 number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
25 protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:8;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:7.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829; the nucleotide sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829; the nucleotide sequence of the full-length protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pj193\_5 deposited  
15 under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 119 to amino acid 128 of SEQ ID NO:8.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and

- (ab) the nucleotide sequence of the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 pj193\_5 deposited under accession number ATCC 98825;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated
- 25 according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 86 to
- 30 nucleotide 829. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from

nucleotide 149 to nucleotide 829, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone

10

pj193\_5 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably

15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 119 to amino acid 128 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an

20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

25

30

- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- 5 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 10 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:9.
- 15 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292; the nucleotide sequence of the full-length protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825. In other preferred embodiments, the
- 20 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most
- 25 preferably thirty) contiguous amino acids of SEQ ID NO:10, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 181 to amino acid 190 of SEQ ID NO:10.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
30 ID NO:9.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:



(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(ab) the nucleotide sequence of the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(bb) the nucleotide sequence of the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:9 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292, to a nucleotide sequence

corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
- (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone

10 pj317\_2 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably

15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 181 to amino acid 190 of SEQ ID NO:10.

In one embodiment, the present invention provides a composition comprising an

20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11;
  - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517;
  - (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 25 NO:11 from nucleotide 904 to nucleotide 2517;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;
  - (e) a polynucleotide encoding the full-length protein encoded by the
- 30 cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:11.
- 15

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517; the nucleotide sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517; the nucleotide sequence of the full-length protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825. In further preferred

20

25 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having

30 biological activity, the fragment comprising the amino acid sequence from amino acid 413 to amino acid 422 of SEQ ID NO:12.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:11.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

5 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

10 (ab) the nucleotide sequence of the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

15 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(bb) the nucleotide sequence of the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:11 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide

2517, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517. Also preferably the polynucleotide isolated according to the above  
5 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
- (b) a fragment of the amino acid sequence of SEQ ID NO:12, the  
15 fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence  
25 from amino acid 413 to amino acid 422 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:13 from nucleotide 18 to nucleotide 257;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:13.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257; the nucleotide sequence of the full-length protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having

biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:14.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:13.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
10 consisting of:

(aa) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and

(ab) the nucleotide sequence of the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and

(bb) the nucleotide sequence of the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:13 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 5 257, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257.

In other embodiments, the present invention provides a composition comprising  
10 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
- (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
- 15 (c) the amino acid sequence encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:14. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:14.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:15 from nucleotide 21 to nucleotide 2432;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825;



- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825;
- 5 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- 10 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 15 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:15.
- 20

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432; the nucleotide sequence of the full-length protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having

25

30

biological activity, the fragment comprising the amino acid sequence from amino acid 397 to amino acid 406 of SEQ ID NO:16.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
10 consisting of:

(aa) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(ab) the nucleotide sequence of the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(bb) the nucleotide sequence of the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
- (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 397 to amino acid 406 of SEQ ID NO:16.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:17.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105; the nucleotide sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105; the nucleotide sequence of the full-length protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 289  
5 to amino acid 298 of SEQ ID NO:18.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:17, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:17; and
    - (ab) the nucleotide sequence of the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
25 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:17, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:17; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 rb649\_3 deposited under accession number ATCC 98825;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:17 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from  
15 nucleotide 501 to nucleotide 2105, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:18;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:18. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:18, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 289 to amino acid 298 of SEQ ID NO:18.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467;
- 5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- 10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- 15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:19.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467; the nucleotide sequence of the full-length protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

30

of clone ca106\_19x deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
5 preferably thirty) contiguous amino acids of SEQ ID NO:20, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 321 to amino acid 330 of SEQ ID NO:20.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
10 ID NO:19.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:  
(i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:19, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:19; and

(ab) the nucleotide sequence of the cDNA insert of clone  
20 ca106\_19x deposited under accession number ATCC 98835;

(ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the  
probe(s);

25 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

(ba) SEQ ID NO:19, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:19; and

(bb) the nucleotide sequence of the cDNA insert of clone  
ca106\_19x deposited under accession number ATCC 98835;



- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:19 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19. Also preferably the  
10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide  
15 509 to nucleotide 2467.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
- 20 (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;

the protein being substantially free from other mammalian proteins. Preferably such  
25 protein comprises the amino acid sequence of SEQ ID NO:20. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a protein comprising a fragment of the amino acid sequence of SEQ  
30 ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 321 to amino acid 330 of SEQ ID NO:20.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least
- 30 25% of the length of SEQ ID NO:21.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802; the nucleotide sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802; the nucleotide sequence of the full-length protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835; or the

nucleotide sequence of a mature protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:22.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:21.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

- (ab) the nucleotide sequence of the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;

- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

- (bb) the nucleotide sequence of the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:21 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
- (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:22.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- 25 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:23.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714; the nucleotide sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714; the nucleotide sequence of the full-length protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 106 to amino acid 115 of SEQ ID NO:24.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:23.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and
    - (ab) the nucleotide sequence of the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide sequences that

**COMPOSITIONS AND METHODS FOR MULTIVALENT, MULTISPECIFIC AGENTS GENERATED FROM  $V_H$  AND  $V_L$  DOMAINS**

Inventor #1: Edmund A. Rossi

Citizenship:

Address:

Inventor #2: Soumitra K. Saha

Citizenship:

Address:

Inventor #3: Chien-Hsing Chang

Citizenship:

Address:



ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013;
- 10          (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- 15          (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- 20          (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- 25          (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30          (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:3.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013; the nucleotide sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013; the nucleotide sequence of the full-length protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825; or the  
5 nucleotide sequence of a mature protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825. In further preferred  
10 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having  
15 biological activity, the fragment comprising the amino acid sequence from amino acid 162 to amino acid 171 of SEQ ID NO:4.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:3.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 25 (aa) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and
- (ab) the nucleotide sequence of the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bb) the nucleotide sequence of the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:3 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of SEQ  
ID NO:4 having biological activity, the fragment comprising the amino acid sequence from  
10 amino acid 162 to amino acid 171 of SEQ ID NO:4.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825;
- 25 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:5.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825; the nucleotide sequence of the full-length  
10 protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825. In further preferred  
15 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having  
20 biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:6.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:5.

Further embodiments of the invention provide isolated polynucleotides produced  
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(ab) the nucleotide sequence of the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(bb) the nucleotide sequence of the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5, and extending  
20 contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:5 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825, and extending  
25 contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising the amino acid sequence from  
10 amino acid 74 to amino acid 83 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pj193\_5 deposited under accession  
20 number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
25 protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:8;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:7.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829; the nucleotide sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829; the nucleotide sequence of the full-length protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pj193\_5 deposited  
15 under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 119 to amino acid 128 of SEQ ID NO:8.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and



- (ab) the nucleotide sequence of the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 pj193\_5 deposited under accession number ATCC 98825;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated
- 25 according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 86 to
- 30 nucleotide 829. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from

nucleotide 149 to nucleotide 829, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone

10 pj193\_5 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably

15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 119 to amino acid 128 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an

20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292;

25 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

30 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825;

- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- 5 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 10 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:9.
- 15 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292; the nucleotide sequence of the full-length protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825. In other preferred embodiments, the
- 20 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a polynucleotide encoding
- 25 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 181 to amino acid 190 of SEQ ID NO:10.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ

30 ID NO:9.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(ab) the nucleotide sequence of the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(bb) the nucleotide sequence of the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:9 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292, to a nucleotide sequence

corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
- (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
10 pj317\_2 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably  
15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 181 to amino acid 190 of SEQ ID NO:10.

In one embodiment, the present invention provides a composition comprising an  
20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
25 NO:11 from nucleotide 904 to nucleotide 2517;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the  
30 cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any  
15 one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:11.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517; the nucleotide sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517; the nucleotide sequence of the full-length protein  
20 coding sequence of clone pt332\_1 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825. In further preferred  
25 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having  
30 biological activity, the fragment comprising the amino acid sequence from amino acid 413 to amino acid 422 of SEQ ID NO:12.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:11.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(ab) the nucleotide sequence of the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(bb) the nucleotide sequence of the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:11 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide

2517, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517. Also preferably the polynucleotide isolated according to the above  
5 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
- (b) a fragment of the amino acid sequence of SEQ ID NO:12, the  
15 fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence  
25 from amino acid 413 to amino acid 422 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:13 from nucleotide 18 to nucleotide 257;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825;



- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:13.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257; the nucleotide sequence of the full-length protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having

biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:14.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:13.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:13, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:13; and

(ab) the nucleotide sequence of the cDNA insert of clone  
qc297\_15 deposited under accession number ATCC 98825;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:13, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:13; and

(bb) the nucleotide sequence of the cDNA insert of clone  
qc297\_15 deposited under accession number ATCC 98825;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:13 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
- (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:14. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:14.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:15.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432; the nucleotide sequence of the full-length protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having

biological activity, the fragment comprising the amino acid sequence from amino acid 397 to amino acid 406 of SEQ ID NO:16.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(ab) the nucleotide sequence of the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(bb) the nucleotide sequence of the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
- (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 397 to amino acid 406 of SEQ ID NO:16.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (e) a polynucleotide encoding the full-length protein encoded by the  
5 cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
10 insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment  
15 comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
20 one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:17.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105; the nucleotide sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105; the nucleotide sequence of the full-length protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825; or the nucleotide sequence of a mature protein coding sequence of clone rb649\_3 deposited  
30 under accession number ATCC 98825. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 289  
5 to amino acid 298 of SEQ ID NO:18.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:17, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:17; and
    - (ab) the nucleotide sequence of the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that  
25 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:17, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:17; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 rb649\_3 deposited under accession number ATCC 98825;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).



Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:17 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from  
15 nucleotide 501 to nucleotide 2105, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:18;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:18. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:18, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 289 to amino acid 298 of SEQ ID NO:18.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467;
- 5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- 10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- 15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:19.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467; the nucleotide sequence of the full-length protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

of clone ca106\_19x deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a polynucleotide encoding  
5 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 321 to amino acid 330 of SEQ ID NO:20.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
10 ID NO:19.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
20 ca106\_19x deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 25 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and
    - (bb) the nucleotide sequence of the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:19 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide
- 15 509 to nucleotide 2467.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
  - 20 (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins. Preferably such
- 25 protein comprises the amino acid sequence of SEQ ID NO:20. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a protein comprising a fragment of the amino acid sequence of SEQ
- 30 ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 321 to amino acid 330 of SEQ ID NO:20.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:21.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802; the nucleotide sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802; the nucleotide sequence of the full-length protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835; or the

nucleotide sequence of a mature protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:22.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:21.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

(ab) the nucleotide sequence of the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

- (bb) the nucleotide sequence of the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:21 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 25 (a) the amino acid sequence of SEQ ID NO:22;
- (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- 30

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:22.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- 25 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and



(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:23.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714; the nucleotide sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714; the nucleotide sequence of the full-length protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 106 to amino acid 115 of SEQ ID NO:24.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:23.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and
    - (ab) the nucleotide sequence of the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and

(bb) the nucleotide sequence of the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:23 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:24;

- (b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and
- (c) the amino acid sequence encoded by the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:24. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:24, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 106 to amino acid 115 of SEQ ID NO:24.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 92 to nucleotide 1726;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:25 from nucleotide 1211 to nucleotide 1726;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pk366\_7 deposited under accession number ATCC 98835;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pk366\_7 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone pk366\_7 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:25.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:25 from nucleotide 92 to nucleotide 1726; the nucleotide sequence of SEQ ID NO:25 from nucleotide 1211 to nucleotide 1726; the nucleotide sequence of the full-length protein coding sequence of clone pk366\_7 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone pk366\_7 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 267 to amino acid 276 of SEQ ID NO:26.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:25.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(ab) the nucleotide sequence of the cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(bb) the nucleotide sequence of the cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:25 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide 92 to nucleotide 1726, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from nucleotide 92 to nucleotide 1726, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide

92 to nucleotide 1726. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide 1211 to nucleotide 1726, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from  
5 nucleotide 1211 to nucleotide 1726, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide 1211 to nucleotide 1726.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:26;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:26. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:26, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 267 to amino acid 276 of SEQ ID NO:26.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:27 from nucleotide 61 to nucleotide 1788;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pl741\_5 deposited under accession number ATCC 98835;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pl741\_5 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:27.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788; the nucleotide sequence of SEQ ID NO:27 from nucleotide 61 to nucleotide 1788; the nucleotide sequence of the full-length protein coding sequence of clone pl741\_5 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone pl741\_5 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having

biological activity, the fragment comprising the amino acid sequence from amino acid 290 to amino acid 299 of SEQ ID NO:28.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:27.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

(ab) the nucleotide sequence of the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

(bb) the nucleotide sequence of the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ



ID NO:27 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 61 to nucleotide 1788, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 61 to nucleotide 1788, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 61 to nucleotide 1788.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
- (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pI741\_5 deposited under accession number ATCC 98835;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 290 to amino acid 299 of SEQ ID NO:28.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pp314\_19 deposited under accession number ATCC 98835;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pp314\_19 deposited under accession number ATCC 98835;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:29.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338; the nucleotide sequence of the full-length protein coding sequence of clone pp314\_19 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone pp314\_19 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 280 to amino acid 289 of SEQ ID NO:30.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:29.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and

(ab) the nucleotide sequence of the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and

(bb) the nucleotide sequence of the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:30. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 280 to amino acid 289 of SEQ ID NO:30.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pv35\_1 deposited under accession number ATCC 98835;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pv35\_1 deposited under accession number ATCC 98835;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:31.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102; the nucleotide sequence of the full-length protein coding sequence of clone pv35\_1 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone pv35\_1 deposited under accession number ATCC 98835. In other preferred embodiments, the
- 30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:32, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 152 to amino acid 161 of SEQ ID NO:32.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:31.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
    - (bb) the nucleotide sequence of the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
  - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:31 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:31, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:31. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:32;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:32. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:32, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 152 to amino acid 161 of SEQ ID NO:32.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 413 to nucleotide 733;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pw337\_6 deposited under accession number ATCC 98835;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pw337\_6 deposited under accession number ATCC 98835;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:33.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:33 from nucleotide 413 to nucleotide 733; the nucleotide sequence of the full-length protein coding sequence of clone pw337\_6 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone pw337\_6 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most



preferably thirty) contiguous amino acids of SEQ ID NO:34, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:34.

5        Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:33.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10        (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and
    - 15        (ab) the nucleotide sequence of the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
  - 20        and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25        (ba) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and
    - (bb) the nucleotide sequence of the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835;
  - 30        (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:33 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:33, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:33. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide 413 to nucleotide 733, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 413 to nucleotide 733, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide 413 to nucleotide 733.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:34;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pw337\_6 deposited under accession number ATCC 98835;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:34. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:34, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:34.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rd610\_1 deposited under accession number ATCC 98835;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rd610\_1 deposited under accession number ATCC 98835;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:35.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938; the nucleotide sequence of the full-length protein coding sequence of clone rd610\_1 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone rd610\_1 deposited under accession number ATCC 98835. In other preferred embodiments, the
- 30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:36, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:36.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:35.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
  - 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and
    - (bb) the nucleotide sequence of the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
  - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:35 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:35, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:35. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:36;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:36. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:36, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:36.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 75 to nucleotide 494;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 447 to nucleotide 494;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rd810\_6 deposited under accession number ATCC 98835;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rd810\_6 deposited under accession number ATCC 98835;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:37.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:37 from nucleotide 75 to nucleotide 494; the nucleotide sequence of SEQ ID NO:37 from nucleotide 447 to nucleotide 494; the nucleotide sequence of the full-length protein coding sequence of clone rd810\_6 deposited under accession number ATCC 98835; or the nucleotide sequence of a mature protein coding sequence of clone rd810\_6 deposited under accession number ATCC 98835. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835. In further preferred

embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a polynucleotide encoding  
5 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:38.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:37.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:37, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:37; and

(ab) the nucleotide sequence of the cDNA insert of clone  
rd810\_6 deposited under accession number ATCC 98835;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the  
probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

30 (ba) SEQ ID NO:37, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:37; and

(bb) the nucleotide sequence of the cDNA insert of clone  
rd810\_6 deposited under accession number ATCC 98835;

(ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:37 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide  
10 494, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 75 to nucleotide 494, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 75 to nucleotide 494. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:37 from nucleotide 447 to nucleotide 494, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 447 to nucleotide 494, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 447 to nucleotide 494.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
- (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and  
25 (c) the amino acid sequence encoded by the cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:38. In further preferred embodiments, the present invention provides a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:38.



In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cf85\_1 deposited under accession number ATCC 98850;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cf85\_1 deposited under accession number ATCC 98850;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
20 amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein  
25 of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:39.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080; the nucleotide sequence of the full-length protein coding sequence of clone cf85\_1 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone cf85\_1 deposited

WO 00/09552

under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 145 to amino acid 154 of SEQ ID NO:40.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:39.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
    - (ab) the nucleotide sequence of the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and

- (bb) the nucleotide sequence of the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:39 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
- (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 145 to amino acid 154 of SEQ ID NO:40.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 161 to nucleotide 1348;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 599 to nucleotide 1348;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:41.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:41 from nucleotide 161 to nucleotide 1348; the nucleotide sequence of SEQ ID NO:41

from nucleotide 599 to nucleotide 1348; the nucleotide sequence of the full-length protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 193 to amino acid 202 of SEQ ID NO:42.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:41.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and
    - (ab) the nucleotide sequence of the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and
- (bb) the nucleotide sequence of the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- 5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
10 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:41 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 161 to nucleotide 1348, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 161 to nucleotide 1348, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 161 to nucleotide 1348. Also preferably the polynucleotide isolated according to the above  
20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 599 to nucleotide 1348, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 599 to nucleotide 1348, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 599 to nucleotide 1348.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
- (b) a fragment of the amino acid sequence of SEQ ID NO:42, the  
30 fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:42. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 193 to amino acid 202 of SEQ ID NO:42.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43 from nucleotide 70 to nucleotide 1386;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone np26\_3 deposited under accession number ATCC 98850;
- 15 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone np26\_3 deposited under accession number ATCC 98850;
- 20 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:44;
- 25 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 30 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

WO 00/09552

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:43.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:43 from nucleotide 70 to nucleotide 1386; the nucleotide sequence of the full-length protein coding sequence of clone np26\_3 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone np26\_3 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence from amino acid 214 to amino acid 223 of SEQ ID NO:44.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:43.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

(ab) the nucleotide sequence of the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and



(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

(bb) the nucleotide sequence of the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:43 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43 from nucleotide 70 to nucleotide 1386, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:43 from nucleotide 70 to nucleotide 1386, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:43 from nucleotide 70 to nucleotide 1386.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:44. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence  
5 from amino acid 214 to amino acid 223 of SEQ ID NO:44.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45;
- 10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide 3515;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm412\_12 deposited under accession number ATCC 98850;
- 15 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm412\_12 deposited under accession number ATCC 98850;
- 20 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
25 amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein  
30 of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:45.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide 3515; the nucleotide sequence of the full-length protein coding sequence of clone pm412\_12 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone pm412\_12 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850. In further preferred  
5  
10  
15  
embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 571 to amino acid 580 of SEQ ID NO:46.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:45.

20 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
25 consisting of:

(aa) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

(ab) the nucleotide sequence of the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;

30 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

(bb) the nucleotide sequence of the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:45 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide  
20 3515, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide 3515, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide 3515.

In other embodiments, the present invention provides a composition comprising  
25 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:46;

(b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and

30 (c) the amino acid sequence encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:46. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence  
5 from amino acid 571 to amino acid 580 of SEQ ID NO:46.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
- 10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:47.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780; the nucleotide sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780; the nucleotide sequence of the full-length protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:48.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:47.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and
    - (ab) the nucleotide sequence of the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and

(bb) the nucleotide sequence of the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:48;

- (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49 from nucleotide 64 to nucleotide 486;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:49 from nucleotide 217 to nucleotide 486;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone pv6\_1 deposited under accession number ATCC 98850;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:50;



- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:49.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:49 from nucleotide 64 to nucleotide 486; the nucleotide sequence of SEQ ID NO:49 from nucleotide 217 to nucleotide 486; the nucleotide sequence of the full-length protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:50.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:49.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and

(ab) the nucleotide sequence of the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and

(bb) the nucleotide sequence of the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:49 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 64 to nucleotide 486, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from nucleotide 64 to nucleotide 486, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide

64 to nucleotide 486. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 217 to nucleotide 486, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from  
5 nucleotide 217 to nucleotide 486, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide 217 to nucleotide 486.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:50;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:50. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:50, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:50.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 379 to nucleotide 3783;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:51 from nucleotide 460 to nucleotide 3783;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;
- 15 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:51.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:51 from nucleotide 379 to nucleotide 3783; the nucleotide sequence of SEQ ID NO:51 from nucleotide 460 to nucleotide 3783; the nucleotide sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938; the nucleotide sequence of the full-length protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850; or the
- 30 nucleotide sequence of a mature protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino

acid sequence of SEQ ID NO:52 from amino acid 536 to amino acid 1135. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, 5 most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 562 to amino acid 571 of SEQ ID NO:52.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ 10 ID NO:51.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize 15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and
    - (ab) the nucleotide sequence of the cDNA insert of clone 20 qs14\_3 deposited under accession number ATCC 98850;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 25 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and
    - (bb) the nucleotide sequence of the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:51 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 379 to nucleotide 3783, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 379 to nucleotide 3783, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide
- 15 379 to nucleotide 3783. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 460 to nucleotide 3783, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 460 to nucleotide 3783, to a nucleotide sequence corresponding to the 3' end
- 20 of said sequence of SEQ ID NO:51 from nucleotide 460 to nucleotide 3783. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938,
- 25 to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30
- (a) the amino acid sequence of SEQ ID NO:52;
  - (b) the amino acid sequence of SEQ ID NO:52 from amino acid 536 to amino acid 1135;
  - (c) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and

(d) the amino acid sequence encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:52 or the amino acid sequence of SEQ ID NO:52 from amino acid 536 to amino acid 1135. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 562 to amino acid 571 of SEQ ID NO:52.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 1 to nucleotide 843;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 469 to nucleotide 843;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qy338\_9 deposited under accession number ATCC 98850;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qy338\_9 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:54;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:53.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:53 from nucleotide 1 to nucleotide 843; the nucleotide sequence of SEQ ID NO:53 from nucleotide 469 to nucleotide 843; the nucleotide sequence of the full-length protein coding sequence of clone qy338\_9 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone qy338\_9 deposited  
15 under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 135 to amino acid 144 of SEQ ID NO:54.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:53.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and



- (ab) the nucleotide sequence of the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 qy338\_9 deposited under accession number ATCC 98850;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:53 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 1 to nucleotide 843, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 1 to nucleotide 843, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide
- 30 1 to nucleotide 843. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 469 to nucleotide 843, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from

nucleotide 469 to nucleotide 843, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 469 to nucleotide 843.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:54;
- (b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
10 qy338\_9 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:54. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably  
15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 135 to amino acid 144 of SEQ ID NO:54.

In one embodiment, the present invention provides a composition comprising an  
20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide 906;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
25 NO:55 from nucleotide 325 to nucleotide 906;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850;
- (e) a polynucleotide encoding the full-length protein encoded by the  
30 cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:55.
- 15

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide 906; the nucleotide sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 906; the nucleotide sequence of the full-length protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:56.

20

25

30

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:55.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

5 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and

10 (ab) the nucleotide sequence of the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

15 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and

(bb) the nucleotide sequence of the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;

25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55, and  
30 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide

906, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide 906, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide 906. Also preferably the polynucleotide isolated according to the above  
5 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 906, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 906, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 906.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;
- (b) a fragment of the amino acid sequence of SEQ ID NO:56, the  
15 fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
- (c) the amino acid sequence encoded by the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence  
25 from amino acid 99 to amino acid 108 of SEQ ID NO:56.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:57 from nucleotide 56 to nucleotide 973;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rd232\_5 deposited under accession number ATCC 98850;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rd232\_5 deposited under accession number ATCC 98850;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:57.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:57 from nucleotide 56 to nucleotide 973; the nucleotide sequence of the full-length protein coding sequence of clone rd232\_5 deposited under accession number ATCC 98850; or the nucleotide sequence of a mature protein coding sequence of clone rd232\_5 deposited under accession number ATCC 98850. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having

biological activity, the fragment comprising the amino acid sequence from amino acid 148 to amino acid 157 of SEQ ID NO:58.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:57.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10

(aa) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(ab) the nucleotide sequence of the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;

15

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25

(ba) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(bb) the nucleotide sequence of the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:57 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 56 to nucleotide 5 973, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 56 to nucleotide 973, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide 56 to nucleotide 973.

In other embodiments, the present invention provides a composition comprising  
10 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;
- (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
- 15 (c) the amino acid sequence encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:58. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 148 to amino acid 157 of SEQ ID NO:58.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:59 from nucleotide 893 to nucleotide 2596;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ck213\_12 deposited under accession number ATCC 98918;



- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ck213\_12 deposited under accession number ATCC 98918;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:59.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:59 from nucleotide 893 to nucleotide 2596; the nucleotide sequence of the full-length protein coding sequence of clone ck213\_12 deposited under accession number ATCC 98918; or the nucleotide sequence of a mature protein coding sequence of clone ck213\_12 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having

biological activity, the fragment comprising the amino acid sequence from amino acid 279 to amino acid 288 of SEQ ID NO:60.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:59.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:59, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:59; and

(ab) the nucleotide sequence of the cDNA insert of clone  
ck213\_12 deposited under accession number ATCC 98918;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

25 (ba) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and

(bb) the nucleotide sequence of the cDNA insert of clone  
ck213\_12 deposited under accession number ATCC 98918;

(ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:59 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 893 to nucleotide 2596, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 893 to nucleotide 2596, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:59 from nucleotide 893 to nucleotide 2596.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
- (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:60. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 279 to amino acid 288 of SEQ ID NO:60.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pg195\_1 deposited under accession number ATCC 98918;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pg195\_1 deposited under accession number ATCC 98918;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:62;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:61.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750; the nucleotide sequence of the full-length protein coding sequence of clone pg195\_1 deposited under accession number ATCC 98918; or the nucleotide sequence of a mature protein coding sequence of clone pg195\_1 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having

biological activity, the fragment comprising the amino acid sequence from amino acid 282 to amino acid 291 of SEQ ID NO:62.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:61.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
10 consisting of:

(aa) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

(ab) the nucleotide sequence of the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

25 (bb) the nucleotide sequence of the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:61 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
- (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:62. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 282 to amino acid 291 of SEQ ID NO:62.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 1234 to nucleotide 1440;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:63.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440; the nucleotide sequence of SEQ ID NO:63 from nucleotide 1234 to nucleotide 1440; the nucleotide sequence of the full-length protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918; or the nucleotide sequence of a mature protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:64, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 44  
5 to amino acid 53 of SEQ ID NO:64.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:63.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:63, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:63; and
    - (ab) the nucleotide sequence of the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that  
25 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:63, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:63; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 pw460\_5 deposited under accession number ATCC 98918;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).



Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:63 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:63, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:63. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 1234 to nucleotide 1440, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from  
15 nucleotide 1234 to nucleotide 1440, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 1234 to nucleotide 1440.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:64;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:64. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:64, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 44 to amino acid 53 of SEQ ID NO:64.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 127 to nucleotide 1356;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least
- 30 25% of the length of SEQ ID NO:65.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356; the nucleotide sequence of SEQ ID NO:65 from nucleotide 127 to nucleotide 1356; the nucleotide sequence of the full-length protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918; or the

nucleotide sequence of a mature protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918. In further preferred  
5 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having  
10 biological activity, the fragment comprising the amino acid sequence from amino acid 213 to amino acid 222 of SEQ ID NO:66.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:65.

Further embodiments of the invention provide isolated polynucleotides produced  
15 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:  
20 (aa) SEQ ID NO:65; and  
(ab) the nucleotide sequence of the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and  
25 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:65; and
    - (bb) the nucleotide sequence of the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:65 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:65. Also preferably the polynucleotide isolated according to the above process comprises a
- 10 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356. Also preferably the polynucleotide
- 15 isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 127 to nucleotide 1356, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 127 to nucleotide 1356, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide
- 20 127 to nucleotide 1356.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;

the protein being substantially free from other mammalian proteins. Preferably such

30 protein comprises the amino acid sequence of SEQ ID NO:66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 213 to amino acid 222 of SEQ ID NO:66.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67;
- (b)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624;
- (c)    a polynucleotide comprising the nucleotide sequence of SEQ ID  
10           NO:67 from nucleotide 542 to nucleotide 1624;
- (d)    a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918;
- (e)    a polynucleotide encoding the full-length protein encoded by the  
15           cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;
- (f)    a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918;
- (g)    a polynucleotide encoding a mature protein encoded by the cDNA  
20           insert of clone qy1261\_2 deposited under accession number ATCC 98918;
- (h)    a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i)    a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;
- 25           (j)    a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k)    a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30           (l)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:67.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624; the nucleotide sequence of SEQ ID NO:67 from nucleotide 542 to nucleotide 1624; the nucleotide sequence of the full-length protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918; or  
5 the nucleotide sequence of a mature protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918. In further preferred  
10 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having  
15 biological activity, the fragment comprising the amino acid sequence from amino acid 231 to amino acid 240 of SEQ ID NO:68.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:67.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:67, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:67; and
    - (ab) the nucleotide sequence of the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and  
30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and

(bb) the nucleotide sequence of the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

10 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:67 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 542 to nucleotide 1624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 542 to nucleotide 1624, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 542 to nucleotide 1624.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:68;

(b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and

(c) the amino acid sequence encoded by the cDNA insert of clone  
qy1261\_2 deposited under accession number ATCC 98918;  
the protein being substantially free from other mammalian proteins. Preferably such  
protein comprises the amino acid sequence of SEQ ID NO:68. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the  
amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably  
comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
of SEQ ID NO:68, or a protein comprising a fragment of the amino acid sequence of SEQ  
ID NO:68 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 231 to amino acid 240 of SEQ ID NO:68.

In one embodiment, the present invention provides a composition comprising an  
isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:69;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:69 from nucleotide 1359 to nucleotide 1817;
- (c) a polynucleotide comprising the nucleotide sequence of the full-  
length protein coding sequence of clone rd432\_4 deposited under accession  
number ATCC 98918;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the  
cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918;
- (e) a polynucleotide comprising the nucleotide sequence of a mature  
protein coding sequence of clone rd432\_4 deposited under accession number  
ATCC 98918;
- 25 (f) a polynucleotide encoding a mature protein encoded by the cDNA  
insert of clone rd432\_4 deposited under accession number ATCC 98918;
- (g) a polynucleotide encoding a protein comprising the amino acid  
sequence of SEQ ID NO:70;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:70 having biological activity, the fragment  
comprising eight contiguous amino acids of SEQ ID NO:70;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of  
(a)-(f) above;



(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:69.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:69 from nucleotide 1359 to nucleotide 1817; the nucleotide sequence of the full-length  
10 protein coding sequence of clone rd432\_4 deposited under accession number ATCC 98918; or the nucleotide sequence of a mature protein coding sequence of clone rd432\_4 deposited under accession number ATCC 98918. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918. In further preferred  
15 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having  
20 biological activity, the fragment comprising the amino acid sequence from amino acid 71 to amino acid 80 of SEQ ID NO:70.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:69.

Further embodiments of the invention provide isolated polynucleotides produced  
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and

(ab) the nucleotide sequence of the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and

(bb) the nucleotide sequence of the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:69 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide 1359 to nucleotide 25 1817, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 1359 to nucleotide 1817, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 1359 to nucleotide 1817.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:70;

(b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:70. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 71 to amino acid 80 of SEQ ID NO:70.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 884 to nucleotide 1195;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 947 to nucleotide 1195;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rb789\_14 deposited under accession  
20 number ATCC 207004;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
25 protein coding sequence of clone rb789\_14 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:72;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:71.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:71 from nucleotide 884 to nucleotide 1195; the nucleotide sequence of SEQ ID NO:71 from nucleotide 947 to nucleotide 1195; the nucleotide sequence of the full-length protein coding sequence of clone rb789\_14 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone rb789\_14 deposited  
15 under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:72.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:71.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

- (ab) the nucleotide sequence of the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 rb789\_14 deposited under accession number ATCC 207004;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:71 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 884 to nucleotide 1195, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 884 to nucleotide 1195, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide
- 30 884 to nucleotide 1195. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 947 to nucleotide 1195, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from

nucleotide 947 to nucleotide 1195, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide 947 to nucleotide 1195.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:72;
- (b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and
- (c) the amino acid sequence encoded by the cDNA insert of clone

10 rb789\_14 deposited under accession number ATCC 207004;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:72. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably

15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:72.

In one embodiment, the present invention provides a composition comprising an

20 isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide 374;

25 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yd137\_1 deposited under accession number ATCC 207004;

30 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone yd137\_1 deposited under accession number ATCC 207004;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any  
15 one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:73.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide 374; the nucleotide sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374; the nucleotide sequence of the full-length protein  
20 coding sequence of clone yd137\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yd137\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004. In further preferred  
25 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having  
30 biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:74.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:73.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

5 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

10 (ab) the nucleotide sequence of the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

15 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

(bb) the nucleotide sequence of the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;

25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73, and  
30 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:73 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide



374, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide 374, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide 374. Also preferably the polynucleotide isolated according to the above  
5 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- (b) a fragment of the amino acid sequence of SEQ ID NO:74, the  
15 fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:74. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence  
25 from amino acid 46 to amino acid 55 of SEQ ID NO:74.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:75 from nucleotide 8 to nucleotide 343;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 50 to nucleotide 343;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yd218\_1 deposited under accession number ATCC 207004;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone yd218\_1 deposited under accession number ATCC 207004;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 20 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:75.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 8 to nucleotide 343; the nucleotide sequence of SEQ ID NO:75 from nucleotide 50 to nucleotide 343; the nucleotide sequence of the full-length protein coding sequence of clone yd218\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yd218\_1 deposited
- 30 under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:76.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:75.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 15 (aa) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and
    - (ab) the nucleotide sequence of the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and
    - 30 (bb) the nucleotide sequence of the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:75 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:75, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:75. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 8 to nucleotide 343, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 8 to nucleotide 343, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 8 to nucleotide 343. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 50 to nucleotide 343, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from  
15 nucleotide 50 to nucleotide 343, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 50 to nucleotide 343.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:76;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:76. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 84 to nucleotide 1679;
- 5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ye11\_1 deposited under accession number ATCC 207004;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ye11\_1 deposited under accession number ATCC 207004;
- 10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ye11\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ye11\_1 deposited under accession number ATCC 207004;
- 15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:77.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 84 to nucleotide 1679; the nucleotide sequence of the full-length protein coding sequence of clone ye11\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone ye11\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

30

of clone ye11\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
5 preferably thirty) contiguous amino acids of SEQ ID NO:78, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 261 to amino acid 270 of SEQ ID NO:78.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
10 ID NO:77.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:  
(i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:77, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:77; and

(ab) the nucleotide sequence of the cDNA insert of clone  
20 ye11\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the  
probe(s);

25 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

(ba) SEQ ID NO:77, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:77; and

(bb) the nucleotide sequence of the cDNA insert of clone  
30 ye11\_1 deposited under accession number ATCC 207004;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:77 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 84 to nucleotide 1679, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 84 to nucleotide 1679, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide
- 15 84 to nucleotide 1679.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
  - 20 (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ye11\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins. Preferably such
- 25 protein comprises the amino acid sequence of SEQ ID NO:78. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of SEQ
- 30 ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 261 to amino acid 270 of SEQ ID NO:78.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:79.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646; the nucleotide sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646; the nucleotide sequence of the full-length protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004; or the



nucleotide sequence of a mature protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 257 to amino acid 266 of SEQ ID NO:80.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:79.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and

- (bb) the nucleotide sequence of the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 25 (a) the amino acid sequence of SEQ ID NO:80;
- (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
- 30

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:80. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 257 to amino acid 266 of SEQ ID NO:80.

5        In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81;
- (b)    a polynucleotide comprising the nucleotide sequence of SEQ ID  
10    NO:81 from nucleotide 954 to nucleotide 2423;
- (c)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423;
- (d)    a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ye78\_1 deposited under accession number  
15    ATCC 207004;
- (e)    a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- (f)    a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ye78\_1 deposited under accession number ATCC  
20    207004;
- (g)    a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- (h)    a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:82;
- (i)    a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;
- (j)    a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k)    a polynucleotide which encodes a species homologue of the protein  
30    of (h) or (i) above ;
- (l)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:81.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:81 from nucleotide 954 to nucleotide 2423; the nucleotide sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423; the nucleotide sequence of the full-length protein coding sequence of clone ye78\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone ye78\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:82, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:82.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:81.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and

(bb) the nucleotide sequence of the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:81 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81 from nucleotide 954 to nucleotide 2423, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 954 to nucleotide 2423, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 954 to nucleotide 2423. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:82;

- (b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:82. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:82, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:82.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 176 to nucleotide 1321;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:83 from nucleotide 233 to nucleotide 1321;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone ye90\_1 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:84;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:83.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:83 from nucleotide 176 to nucleotide 1321; the nucleotide sequence of SEQ ID NO:83 from nucleotide 233 to nucleotide 1321; the nucleotide sequence of the full-length protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:84, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 186 to amino acid 195 of SEQ ID NO:84.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:83.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:83, but excluding the poly(A) tail at the 3' end of SEQ ID NO:83; and

(ab) the nucleotide sequence of the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:83, but excluding the poly(A) tail at the 3' end of SEQ ID NO:83; and

(bb) the nucleotide sequence of the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:83 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:83, but excluding the poly(A) tail at the 3' end of SEQ ID NO:83. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 176 to nucleotide 1321, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:83 from nucleotide 176 to nucleotide 1321, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide



176 to nucleotide 1321. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 233 to nucleotide 1321, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:83 from nucleotide 233 to nucleotide 1321, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide 233 to nucleotide 1321.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:84;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:84. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 20 of SEQ ID NO:84, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 186 to amino acid 195 of SEQ ID NO:84.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yi62\_1 deposited under accession number
- 30 ATCC 207004;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;

- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone yi62\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:85.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605; the nucleotide sequence of the full-length protein coding sequence of clone yi62\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yi62\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:86, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:86.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:85.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 10 (aa) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and
    - (ab) the nucleotide sequence of the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 15 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that
  - 20 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and
    - (bb) the nucleotide sequence of the cDNA insert of clone
    - 25 yi62\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:85 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605, to a nucleotide  
5 sequence corresponding to the 3' end of said sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:86;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:86. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:86, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:86.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:87 from nucleotide 430 to nucleotide 798;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:87.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798; the nucleotide sequence of SEQ ID NO:87 from nucleotide 430 to nucleotide 798; the nucleotide sequence of the full-length protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the

fragment comprising the amino acid sequence from amino acid 91 to amino acid 100 of SEQ ID NO:88.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:87.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10

(aa) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and

(ab) the nucleotide sequence of the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;

15

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25

(ba) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and

(bb) the nucleotide sequence of the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:87 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 430 to nucleotide 798, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from nucleotide 430 to nucleotide 798, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 430 to nucleotide 798.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
- (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:88. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 91 to amino acid 100 of SEQ ID NO:88.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 211 to nucleotide 942;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:90;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:89.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:89 from nucleotide 211 to nucleotide 942; the nucleotide sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942; the nucleotide sequence of the full-length protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the

30



polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:90, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:90.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:89.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and
    - (ab) the nucleotide sequence of the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and
    - (bb) the nucleotide sequence of the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:89 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89 from nucleotide 211 to nucleotide 942, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from nucleotide 211 to nucleotide 942, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:89 from nucleotide
- 15 211 to nucleotide 942. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942, to a nucleotide sequence corresponding to the 3' end of
- 20 said sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:90;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;

the protein being substantially free from other mammalian proteins. Preferably such

30 protein comprises the amino acid sequence of SEQ ID NO:90. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:90, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:90.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone yt14\_1 deposited under accession number  
10           ATCC 207004;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;
- (e) a polynucleotide comprising the nucleotide sequence of a mature  
15           protein coding sequence of clone yt14\_1 deposited under accession number ATCC 207004;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;
- (g) a polynucleotide encoding a protein comprising the amino acid  
20           sequence of SEQ ID NO:92;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of  
25           (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30           (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:91.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784; the nucleotide sequence of the full-length

protein coding sequence of clone yt14\_1 deposited under accession number ATCC 207004; or the nucleotide sequence of a mature protein coding sequence of clone yt14\_1 deposited under accession number ATCC 207004. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004. In further preferred  
5 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a polynucleotide encoding  
10 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising the amino acid sequence from amino acid 101 to amino acid 110 of SEQ ID NO:92.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:91.

15 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
20 consisting of:
    - (aa) SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91; and
    - (ab) the nucleotide sequence of the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;
  - 25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- 30 (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91; and

(bb) the nucleotide sequence of the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:91 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:92;

25 (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and

(c) the amino acid sequence encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:92. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:92 having biological activity, the fragment comprising the amino acid sequence from amino acid 101 to amino acid 110 of SEQ ID NO:92.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93;
- (b)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441;
- (c)    a polynucleotide comprising the nucleotide sequence of the full-  
10   length protein coding sequence of clone bf157\_16 deposited under accession number ATCC 207088;
- (d)    a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
- (e)    a polynucleotide comprising the nucleotide sequence of a mature  
15   protein coding sequence of clone bf157\_16 deposited under accession number ATCC 207088;
- (f)    a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
- (g)    a polynucleotide encoding a protein comprising the amino acid  
20   sequence of SEQ ID NO:94;
- (h)    a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (i)    a polynucleotide which is an allelic variant of a polynucleotide of  
25   (a)-(f) above;
- (j)    a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30          (l)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:93.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441; the nucleotide sequence of the full-length

protein coding sequence of clone bf157\_16 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone bf157\_16 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088. In further preferred  
5       embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a polynucleotide encoding  
10       a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 220 to amino acid 229 of SEQ ID NO:94.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:93.

15       Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
20       consisting of:
    - (aa) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and
    - (ab) the nucleotide sequence of the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
  - 25       (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- 30       (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and
- (bb) the nucleotide sequence of the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
- 5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:93 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
- (b) a fragment of the amino acid sequence of SEQ ID NO:94, the  
25 fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:94. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a protein comprising a fragment of the amino acid sequence of SEQ



ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 220 to amino acid 229 of SEQ ID NO:94.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bk343\_2 deposited under accession  
10 number ATCC 207088;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature  
15 protein coding sequence of clone bk343\_2 deposited under accession number ATCC 207088;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid  
20 sequence of SEQ ID NO:96;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of  
25 (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:95.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629; the nucleotide sequence of the full-length

protein coding sequence of clone bk343\_2 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone bk343\_2 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:96, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 63 to amino acid 72 of SEQ ID NO:96.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:95.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and
    - (ab) the nucleotide sequence of the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

(bb) the nucleotide sequence of the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
10 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:95 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:96;

(b) a fragment of the amino acid sequence of SEQ ID NO:96, the  
25 fragment comprising eight contiguous amino acids of SEQ ID NO:96; and

(c) the amino acid sequence encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:96. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:96, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 63 to amino acid 72 of SEQ ID NO:96.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97;
- (b)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97 from nucleotide 556 to nucleotide 951;
- (c)    a polynucleotide comprising the nucleotide sequence of SEQ ID  
10       NO:97 from nucleotide 868 to nucleotide 951;
- (d)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295;
- (e)    a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cd205\_2 deposited under accession  
15       number ATCC 207088;
- (f)    a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
- (g)    a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cd205\_2 deposited under accession number  
20       ATCC 207088;
- (h)    a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
- (i)    a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:98;
- 25       (j)    a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;
- (k)    a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- 30       (l)    a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

(n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:97.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:97 from nucleotide 556 to nucleotide 951; the nucleotide sequence of SEQ ID NO:97 from nucleotide 868 to nucleotide 951; the nucleotide sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295; the nucleotide sequence of the full-length protein coding sequence of clone cd205\_2 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone cd205\_2 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:98, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:98.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:97.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and
    - (ab) the nucleotide sequence of the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and

10 (bb) the nucleotide sequence of the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:97 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:97, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:97. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 556 to nucleotide 951, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 556 to nucleotide 951, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide 556 to nucleotide 951. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 868 to nucleotide 951, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from  
30 nucleotide 868 to nucleotide 951, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide 868 to nucleotide 951. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295, and extending contiguously from a nucleotide sequence corresponding

to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295.

In other embodiments, the present invention provides a composition comprising  
5 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:98;
- (b) a fragment of the amino acid sequence of SEQ ID NO:98, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:98; and
- 10 (c) the amino acid sequence encoded by the cDNA insert of clone  
cd205\_2 deposited under accession number ATCC 207088;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:98. In further preferred  
15 amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:98, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:98.

20 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:99;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
25 NO:99 from nucleotide 216 to nucleotide 443;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:99 from nucleotide 306 to nucleotide 443;
- (d) a polynucleotide comprising the nucleotide sequence of the full-  
length protein coding sequence of clone cw1292\_8 deposited under accession  
30 number ATCC 207088;
- (e) a polynucleotide encoding the full-length protein encoded by the  
cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cw1292\_8 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:99.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:99 from nucleotide 216 to nucleotide 443; the nucleotide sequence of SEQ ID NO:99 from nucleotide 306 to nucleotide 443; the nucleotide sequence of the full-length protein coding sequence of clone cw1292\_8 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone cw1292\_8 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:100.



Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:99.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 10 (aa) SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99; and
    - (ab) the nucleotide sequence of the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 15 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that
  - 20 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99; and
    - (bb) the nucleotide sequence of the cDNA insert of clone
    - 25 cw1292\_8 deposited under accession number ATCC 207088;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:99 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99 from nucleotide 216 to nucleotide 443, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:99 from nucleotide 216 to nucleotide 443, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:99 from nucleotide 216 to nucleotide 443. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99 from nucleotide 306 to nucleotide 443, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:99 from nucleotide 306 to nucleotide 443, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:99 from nucleotide 306 to nucleotide 443.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:100. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:100.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cw1475\_2 deposited under accession number ATCC 207088;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cw1475\_2 deposited under accession number ATCC 207088;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:101.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447; the nucleotide sequence of the full-length protein coding sequence of clone cw1475\_2 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone cw1475\_2 deposited under accession number ATCC 207088. In other preferred
- 30 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:102, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:102.

- 5           Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:101.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10           (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and
    - 15           (ab) the nucleotide sequence of the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the probe(s);
    - 20           and
  - (b) a process comprising the steps of:
    - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - 25           (ba) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and
      - (bb) the nucleotide sequence of the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
      - 30           (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
      - (iii) amplifying human DNA sequences; and
      - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:101 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:101, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:101. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447,  
10 to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:102;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:102. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:102, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:102.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dd428\_4 deposited under accession number ATCC 207088;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dd428\_4 deposited under accession number ATCC 207088;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment
- 15 comprising eight contiguous amino acids of SEQ ID NO:104;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:103.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954; the nucleotide sequence of the full-length protein coding sequence of clone dd428\_4 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone dd428\_4 deposited under accession number ATCC 207088. In other preferred embodiments, the
- 30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:104, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 102 to amino acid 111 of SEQ ID NO:104.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:103.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and
    - (bb) the nucleotide sequence of the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
  - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:103 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:103, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:103. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:104;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:104. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:104, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 102 to amino acid 111 of SEQ ID NO:104.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895;



- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dh1073\_12 deposited under accession number ATCC 207088;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dh1073\_12 deposited under accession number ATCC 207088;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:105.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895; the nucleotide sequence of the full-length protein coding sequence of clone dh1073\_12 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone dh1073\_12 deposited under accession number ATCC 207088. In other preferred
- 30 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088.
- In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 28 to amino acid 37 of SEQ ID NO:106.

- 5           Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:105.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a)   a process comprising the steps of:
- 10           (i)   preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa)   SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and
- 15           (ab)   the nucleotide sequence of the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- (ii)   hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii)   isolating the DNA polynucleotides detected with the
- 20           probe(s);
- and
- (b)   a process comprising the steps of:
- (i)   preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from
- 25           the group consisting of:
- (ba)   SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and
- (bb)   the nucleotide sequence of the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- 30           (ii)   hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii)   amplifying human DNA sequences; and
- (iv)   isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:105 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:105, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895,  
10 to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:106;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:106. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:106, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 28 to amino acid 37 of SEQ ID NO:106.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dw78\_1 deposited under accession number ATCC 207088;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dw78\_1 deposited under accession number ATCC 207088;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:108;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment  
15 comprising eight contiguous amino acids of SEQ ID NO:108;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:107.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791; the nucleotide sequence of the full-length protein coding sequence of clone dw78\_1 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone dw78\_1 deposited under accession number ATCC 207088. In other preferred embodiments, the  
30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:108, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 56 to amino acid 65 of SEQ ID NO:108.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:107.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and
- 15 (ab) the nucleotide sequence of the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- 20 and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 25 (ba) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and
- (bb) the nucleotide sequence of the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:107 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:107, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:107. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:108;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:108. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:108, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 56 to amino acid 65 of SEQ ID NO:108.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fh116\_11 deposited under accession number ATCC 207088;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fh116\_11 deposited under accession number ATCC 207088;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:109.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944; the nucleotide sequence of the full-length protein coding sequence of clone fh116\_11 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone fh116\_11 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:110, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 136 to amino acid 145 of SEQ ID NO:110.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:109.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
  - 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and
    - (bb) the nucleotide sequence of the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
  - 25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - 30 (iv) isolating the polynucleotide products of step (b)(iii).



Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:109 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:109, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:109. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:110;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:110. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:110, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 136 to amino acid 145 of SEQ ID NO:110.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fy356\_14 deposited under accession number ATCC 207088;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fy356\_14 deposited under accession number ATCC 207088;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:111.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610; the nucleotide sequence of the full-length protein coding sequence of clone fy356\_14 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone fy356\_14 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:112, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 238 to amino acid 247 of SEQ ID NO:112.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:111.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
  - 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and
    - (bb) the nucleotide sequence of the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
  - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:111 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:111, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:111. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:112;
- (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
fy356\_14 deposited under accession number ATCC 207088;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:112. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:112, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 238 to amino acid 247 of SEQ ID NO:112.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide 669;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 112 to nucleotide 669;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:114;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:113.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide 669; the nucleotide sequence of SEQ ID NO:113 from nucleotide 112 to nucleotide 669; the nucleotide sequence of the full-length protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088; or the nucleotide sequence of a mature protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088. In further preferred embodiments, the

present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:114, or a polynucleotide encoding a protein comprising a  
5 fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 98 to amino acid 107 of SEQ ID NO:114.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:113.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
15 consisting of:

(aa) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and

(ab) the nucleotide sequence of the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and

(bb) the nucleotide sequence of the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:113 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide  
10 669, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide 669, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide 669. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:113 from nucleotide 112 to nucleotide 669, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 112 to nucleotide 669, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:113 from nucleotide 112 to nucleotide 669.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:114;
- (b) a fragment of the amino acid sequence of SEQ ID NO:114, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:114; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
25 iw66\_1 deposited under accession number ATCC 207088;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:114. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:114, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 98 to amino acid 107 of SEQ ID NO:114.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone kh13\_4 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone kh13\_4 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:115.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416; the nucleotide sequence of the full-length protein coding sequence of clone kh13\_4 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone kh13\_4



deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:116, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 37 to amino acid 46 of SEQ ID NO:116.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:115.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:115, but excluding the poly(A) tail at the  
20 3' end of SEQ ID NO:115; and
    - (ab) the nucleotide sequence of the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115; and

- (bb) the nucleotide sequence of the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:115 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:116;
- (b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and
- (c) the amino acid sequence encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:116. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:116, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:116.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ko258\_4 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ko258\_4 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:117.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602; the nucleotide sequence of the full-length protein coding sequence of clone ko258\_4 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone ko258\_4

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:118, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 61 to amino acid 70 of SEQ ID NO:118.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:117.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:117, but excluding the poly(A) tail at the  
20 3' end of SEQ ID NO:117; and
- (ab) the nucleotide sequence of the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117; and

- (bb) the nucleotide sequence of the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:117 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;
- (b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:118. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:118, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:118.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:119;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:119 from nucleotide 434 to nucleotide 739;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone kv10\_8 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone kv10\_8 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:120;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
20 amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:120;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein  
25 of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any  
30 one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:119.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:119 from nucleotide 434 to nucleotide 739; the nucleotide sequence of the full-length protein coding sequence of clone kv10\_8 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone kv10\_8

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:120, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 46 to amino acid 55 of SEQ ID NO:120.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:119.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 20 (aa) SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119; and
- (ab) the nucleotide sequence of the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119; and

- (bb) the nucleotide sequence of the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:119, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:119 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:119 from nucleotide 434 to nucleotide 739, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:119 from nucleotide 434 to nucleotide 739, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:119 from nucleotide 434 to nucleotide 739.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:120;
- (b) a fragment of the amino acid sequence of SEQ ID NO:120, the fragment comprising eight contiguous amino acids of SEQ ID NO:120; and
- (c) the amino acid sequence encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;

25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:120. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:120, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:120.



In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone LL89\_3 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone LL89\_3 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:122;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:122;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:121.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310; the nucleotide sequence of the full-length protein coding sequence of clone LL89\_3 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone LL89\_3

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:122, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:122.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:121.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121; and
    - (ab) the nucleotide sequence of the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121; and

- (bb) the nucleotide sequence of the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:121, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ  
10 ID NO:121 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
15 of said sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
20 consisting of:

- (a) the amino acid sequence of SEQ ID NO:122;
- (b) a fragment of the amino acid sequence of SEQ ID NO:122, the fragment comprising eight contiguous amino acids of SEQ ID NO:122; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
25 LL89\_3 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:122. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably  
30 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:122, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:122.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:123;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:123 from nucleotide 22 to nucleotide 288;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone mc300\_1 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone mc300\_1 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:124;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:124;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:123.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:123 from nucleotide 22 to nucleotide 288; the nucleotide sequence of the full-length protein coding sequence of clone mc300\_1 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone mc300\_1

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089. In further preferred  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:124, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 39 to amino acid 48 of SEQ ID NO:124.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:123.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:123, but excluding the poly(A) tail at the  
20 3' end of SEQ ID NO:123; and
    - (ab) the nucleotide sequence of the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:123, but excluding the poly(A) tail at the 3' end of SEQ ID NO:123; and

- (bb) the nucleotide sequence of the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:123, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:123 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:123, but excluding the poly(A) tail at the 3' end of SEQ ID NO:123. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:123 from nucleotide 22 to nucleotide 288, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:123 from nucleotide 22 to nucleotide 288, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:123 from nucleotide 22 to nucleotide 288.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:124;
- (b) a fragment of the amino acid sequence of SEQ ID NO:124, the fragment comprising eight contiguous amino acids of SEQ ID NO:124; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:124. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:124, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:124.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ml227\_1 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ml227\_1 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:126;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:126;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:125.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449; the nucleotide sequence of the full-length protein coding sequence of clone ml227\_1 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone ml227\_1

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089. In further preferred  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:126, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 370 to amino acid 379 of SEQ ID NO:126.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:125.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:125, but excluding the poly(A) tail at the  
20 3' end of SEQ ID NO:125; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:125, but excluding the poly(A) tail at the 3' end of SEQ ID NO:125; and



- (bb) the nucleotide sequence of the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ  
10 ID NO:125 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:125, but excluding the poly(A) tail at the 3' end of SEQ ID NO:125. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
15 of said sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
20 consisting of:

- (a) the amino acid sequence of SEQ ID NO:126;
- (b) a fragment of the amino acid sequence of SEQ ID NO:126, the fragment comprising eight contiguous amino acids of SEQ ID NO:126; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
25 ml227\_1 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:126. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment preferably  
30 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:126, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising the amino acid sequence from amino acid 370 to amino acid 379 of SEQ ID NO:126.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127 from nucleotide 82 to nucleotide 1980;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone mm367\_6 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone mm367\_6 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:128;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:128;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:127.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:127 from nucleotide 82 to nucleotide 1980; the nucleotide sequence of the full-length protein coding sequence of clone mm367\_6 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone mm367\_6

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089. In further preferred  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:128, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 311 to amino acid 320 of SEQ ID NO:128.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:127.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:127, but excluding the poly(A) tail at the  
20 3' end of SEQ ID NO:127; and
    - (ab) the nucleotide sequence of the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:127, but excluding the poly(A) tail at the 3' end of SEQ ID NO:127; and

- (bb) the nucleotide sequence of the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:127 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:127, but excluding the poly(A) tail at the 3' end of SEQ ID NO:127. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127 from nucleotide 82 to nucleotide 1980, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:127 from nucleotide 82 to nucleotide 1980, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:127 from nucleotide 82 to nucleotide 1980.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:128;
- (b) a fragment of the amino acid sequence of SEQ ID NO:128, the fragment comprising eight contiguous amino acids of SEQ ID NO:128; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;

25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:128. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:128, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising the amino acid sequence from amino acid 311 to amino acid 320 of SEQ ID NO:128.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone mt124\_3 deposited under accession number ATCC 207089;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone mt124\_3 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:130;
- (h) a polynucleotide encoding a protein comprising a fragment of the  
20 amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:130;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein  
25 of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least  
30 25% of the length of SEQ ID NO:129.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856; the nucleotide sequence of the full-length protein coding sequence of clone mt124\_3 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone mt124\_3

deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:130, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:130.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:129.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:129, but excluding the poly(A) tail at the 3' end of SEQ ID NO:129; and
    - (ab) the nucleotide sequence of the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:129, but excluding the poly(A) tail at the 3' end of SEQ ID NO:129; and

- (bb) the nucleotide sequence of the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:129 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:129, but excluding the poly(A) tail at the 3' end of SEQ ID NO:129. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:130;
- (b) a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;

25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:130. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:130, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:130.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131 from nucleotide 901 to nucleotide 2940;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:132;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:132;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:131.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940; the nucleotide sequence of SEQ ID



NO:131 from nucleotide 901 to nucleotide 2940; the nucleotide sequence of the full-length protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:132, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising the amino acid sequence from amino acid 342 to amino acid 351 of SEQ ID NO:132.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:131.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131; and
    - (ab) the nucleotide sequence of the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131; and

(bb) the nucleotide sequence of the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:131, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:131 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940. Also preferably the polynucleotide isolated according to the above  
20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:131 from nucleotide 901 to nucleotide 2940, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:131 from nucleotide 901 to nucleotide 2940, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:131 from nucleotide 901 to nucleotide 2940.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:132;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:132, the fragment comprising eight contiguous amino acids of SEQ ID NO:132; and

(c) the amino acid sequence encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:132. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:132, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising the amino acid sequence from amino acid 342 to amino acid 351 of SEQ ID NO:132.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:134;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:134;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:133.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448; the nucleotide sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448; the nucleotide sequence of the full-length protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:134, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising the amino acid sequence from amino acid 49 to amino acid 58 of SEQ ID NO:134.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:133.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:133, but excluding the poly(A) tail at the 3' end of SEQ ID NO:133; and

(ab) the nucleotide sequence of the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:133, but excluding the poly(A) tail at the 3' end of SEQ ID NO:133; and

(bb) the nucleotide sequence of the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:133, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:133 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:133, but excluding the poly(A) tail at the 3' end of SEQ ID NO:133. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:134;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:134, the fragment comprising eight contiguous amino acids of SEQ ID NO:134; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
5      qy442\_2 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:134. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably  
10   comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:134, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising the amino acid sequence from amino acid 49 to amino acid 58 of SEQ ID NO:134.

In one embodiment, the present invention provides a composition comprising an  
15   isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:135;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:135 from nucleotide 28 to nucleotide 777;
- 20      (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:135 from nucleotide 73 to nucleotide 777;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089;
- 25      (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089;
- 30      (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:136;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:136;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:135.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:135 from nucleotide 28 to nucleotide 777; the nucleotide sequence of SEQ ID NO:135 from nucleotide 73 to nucleotide 777; the nucleotide sequence of the full-length protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:136, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment comprising the amino acid sequence from amino acid 120 to amino acid 129 of SEQ ID NO:136.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:135.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135; and

(ab) the nucleotide sequence of the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135; and

(bb) the nucleotide sequence of the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:135, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:135 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:135 from nucleotide 28 to nucleotide 777, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:135 from nucleotide 28 to nucleotide 777, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:135 from nucleotide



28 to nucleotide 777. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:135 from nucleotide 73 to nucleotide 777, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:135 from  
5 nucleotide 73 to nucleotide 777, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:135 from nucleotide 73 to nucleotide 777.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:136;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:136, the fragment comprising eight contiguous amino acids of SEQ ID NO:136; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:136. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:136, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment comprising the amino acid sequence from amino acid 120 to amino acid 129 of SEQ ID NO:136.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:137;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:137 from nucleotide 179 to nucleotide 745;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:137 from nucleotide 233 to nucleotide 745;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone rk80\_3 deposited under accession number ATCC 207089;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone rk80\_3 deposited under accession number ATCC 207089;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:138;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:138;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:137.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:137 from nucleotide 179 to nucleotide 745; the nucleotide sequence of SEQ ID NO:137 from nucleotide 233 to nucleotide 745; the nucleotide sequence of the full-length protein coding sequence of clone rk80\_3 deposited under accession number ATCC 207089; or the nucleotide sequence of a mature protein coding sequence of clone rk80\_3 deposited under accession number ATCC 207089. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:138, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the

fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:138.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:137.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:137, but excluding the poly(A) tail at the 3' end of SEQ ID NO:137; and

(ab) the nucleotide sequence of the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:137, but excluding the poly(A) tail at the 3' end of SEQ ID NO:137; and

(bb) the nucleotide sequence of the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:137, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:137 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:137, but excluding the poly(A) tail at the 3' end of SEQ ID NO:137. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:137 from nucleotide 179 to nucleotide 745, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:137 from nucleotide 179 to nucleotide 745, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:137 from nucleotide 179 to nucleotide 745. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:137 from nucleotide 233 to nucleotide 745, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:137 from nucleotide 233 to nucleotide 745, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:137 from nucleotide 233 to nucleotide 745.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:138;
- (b) a fragment of the amino acid sequence of SEQ ID NO:138, the fragment comprising eight contiguous amino acids of SEQ ID NO:138; and
- (c) the amino acid sequence encoded by the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:138. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:138, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:138.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:139;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone au36\_42 deposited under accession number ATCC 207187;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone au36\_42 deposited under accession number ATCC 207187;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:140;
- 15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:140;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 20 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:139.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274; the nucleotide sequence of the full-length protein coding sequence of clone au36\_42 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone au36\_42 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding

30

a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:140, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140  
5 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:140.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:139.

Further embodiments of the invention provide isolated polynucleotides produced  
10 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:139, but excluding the poly(A) tail at the 3' end of SEQ ID NO:139; and

(ab) the nucleotide sequence of the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:139, but excluding the poly(A) tail at the 3' end of SEQ ID NO:139; and

30 (bb) the nucleotide sequence of the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:139, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:139 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:139, but excluding the poly(A) tail at the 3' end of SEQ ID NO:139. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:140;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:140, the fragment comprising eight contiguous amino acids of SEQ ID NO:140; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:140. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:140, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:140.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:141;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bo549\_13 deposited under accession number ATCC 207187;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone bo549\_13 deposited under accession number ATCC 207187;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:142;
- 15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:142;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:141.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774; the nucleotide sequence of the full-length protein coding sequence of clone bo549\_13 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone bo549\_13 deposited under accession number ATCC 207187. In other preferred embodiments, the
- 30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment preferably comprising eight (more preferably twenty, most



preferably thirty) contiguous amino acids of SEQ ID NO:142, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment comprising the amino acid sequence from amino acid 27 to amino acid 36 of SEQ ID NO:142.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:141.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:141, but excluding the poly(A) tail at the 3' end of SEQ ID NO:141; and
    - 15 (ab) the nucleotide sequence of the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
  - 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:141, but excluding the poly(A) tail at the 3' end of SEQ ID NO:141; and
    - (bb) the nucleotide sequence of the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
  - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:141, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:141 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:141, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:141. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:142;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:142, the fragment comprising eight contiguous amino acids of SEQ ID NO:142; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:142. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:142, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment comprising the amino acid sequence from amino acid 27 to amino acid 36 of SEQ ID NO:142.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:143;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide 969;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:143 from nucleotide 385 to nucleotide 969;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:144;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:144;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:143.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide 969; the nucleotide sequence of SEQ ID NO:143 from nucleotide 385 to nucleotide 969; the nucleotide sequence of the full-length protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187. In further preferred

embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:144, or a polynucleotide  
5 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment comprising the amino acid sequence from amino acid 128 to amino acid 137 of SEQ ID NO:144.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:143.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:143, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:143; and

(ab) the nucleotide sequence of the cDNA insert of clone  
da529\_3 deposited under accession number ATCC 207187;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the  
probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

(ba) SEQ ID NO:143, but excluding the poly(A) tail at the  
30 3' end of SEQ ID NO:143; and

(bb) the nucleotide sequence of the cDNA insert of clone  
da529\_3 deposited under accession number ATCC 207187;

(ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:143, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:143 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:143, but excluding the poly(A) tail at the 3' end of SEQ ID NO:143. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide  
10 969, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide 969, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide 969. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:143 from nucleotide 385 to nucleotide 969, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:143 from nucleotide 385 to nucleotide 969, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:143 from nucleotide 385 to nucleotide 969.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:144;
- (b) a fragment of the amino acid sequence of SEQ ID NO:144, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:144; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:144. In further preferred  
embodiments, the present invention provides a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:144 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:144, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment comprising the amino acid sequence from amino acid 128 to amino acid 137 of SEQ ID NO:144.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:145;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:145 from nucleotide 368 to nucleotide 667;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:146;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:146;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:145.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667; the nucleotide sequence of SEQ ID NO:145

from nucleotide 368 to nucleotide 667; the nucleotide sequence of the full-length protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:146, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:146.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:145.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:145, but excluding the poly(A) tail at the 3' end of SEQ ID NO:145; and
    - (ab) the nucleotide sequence of the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:145, but excluding the poly(A) tail at the 3' end of SEQ ID NO:145; and

(bb) the nucleotide sequence of the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:145, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:145 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:145, but excluding the poly(A) tail at the 3' end of SEQ ID NO:145. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence

15 corresponding to the cDNA sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667. Also preferably the polynucleotide isolated according to the above

20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:145 from nucleotide 368 to nucleotide 667, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:145 from nucleotide 368 to nucleotide 667, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:145 from nucleotide 368 to nucleotide 667.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:146;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:146, the fragment comprising eight contiguous amino acids of SEQ ID NO:146; and

(c) the amino acid sequence encoded by the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:146. In further preferred



embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:146, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:146.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:147;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:147 from nucleotide 103 to nucleotide 1368;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fa171\_1 deposited under accession number ATCC 207187;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fa171\_1 deposited under accession number ATCC 207187;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:148;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:148;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:147.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID  
5 NO:147 from nucleotide 103 to nucleotide 1368; the nucleotide sequence of the full-length  
protein coding sequence of clone fa171\_1 deposited under accession number ATCC  
207187; or the nucleotide sequence of a mature protein coding sequence of clone fa171\_1  
deposited under accession number ATCC 207187. In other preferred embodiments, the  
polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
10 of clone fa171\_1 deposited under accession number ATCC 207187. In further preferred  
embodiments, the present invention provides a polynucleotide encoding a protein  
comprising a fragment of the amino acid sequence of SEQ ID NO:148 having biological  
activity, the fragment preferably comprising eight (more preferably twenty, most  
preferably thirty) contiguous amino acids of SEQ ID NO:148, or a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148  
having biological activity, the fragment comprising the amino acid sequence from amino  
acid 206 to amino acid 215 of SEQ ID NO:148.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
ID NO:147.

20 Further embodiments of the invention provide isolated polynucleotides produced  
according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
25 consisting of:
    - (aa) SEQ ID NO:147, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:147; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
fa171\_1 deposited under accession number ATCC 207187;
  - 30 (ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the  
probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:147, but excluding the poly(A) tail at the 3' end of SEQ ID NO:147; and

(bb) the nucleotide sequence of the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:147, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:147 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:147, but excluding the poly(A) tail at the 3' end of SEQ ID NO:147. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:147 from nucleotide  
20 1368, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:147 from nucleotide 103 to nucleotide 1368, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:147 from nucleotide 103 to nucleotide 1368.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:148;

(b) a fragment of the amino acid sequence of SEQ ID NO:148, the fragment comprising eight contiguous amino acids of SEQ ID NO:148; and

30 (c) the amino acid sequence encoded by the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:148. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:148 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:148, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148 having biological activity, the fragment comprising the amino acid sequence  
5 from amino acid 206 to amino acid 215 of SEQ ID NO:148.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149;
- 10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407;
- (d) a polynucleotide comprising the nucleotide sequence of the full-  
15 length protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20 protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25 sequence of SEQ ID NO:150;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:150;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30 (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:149.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407; the nucleotide sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407; the nucleotide sequence of the full-length protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:150, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment comprising the amino acid sequence from amino acid 198 to amino acid 207 of SEQ ID NO:150.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:149.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:149, but excluding the poly(A) tail at the 3' end of SEQ ID NO:149; and
    - (ab) the nucleotide sequence of the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:149, but excluding the poly(A) tail at the 3' end of SEQ ID NO:149; and

(bb) the nucleotide sequence of the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:149, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:149 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:149, but excluding the poly(A) tail at the 3' end of SEQ ID NO:149. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:150;

- (b) a fragment of the amino acid sequence of SEQ ID NO:150, the fragment comprising eight contiguous amino acids of SEQ ID NO:150; and
- (c) the amino acid sequence encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:150. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:150, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment comprising the amino acid sequence from amino acid 198 to amino acid 207 of SEQ ID NO:150.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:151;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:151 from nucleotide 301 to nucleotide 1035;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:151 from nucleotide 916 to nucleotide 1035;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone pe246\_1 deposited under accession number ATCC 207187;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:152;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:152;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:151.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:151 from nucleotide 301 to nucleotide 1035; the nucleotide sequence of SEQ ID NO:151 from nucleotide 916 to nucleotide 1035; the nucleotide sequence of the full-length protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:152, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:152.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:151.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:



(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:151, but excluding the poly(A) tail at the 3' end of SEQ ID NO:151; and

(ab) the nucleotide sequence of the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:151, but excluding the poly(A) tail at the 3' end of SEQ ID NO:151; and

(bb) the nucleotide sequence of the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:151, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:151 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:151, but excluding the poly(A) tail at the 3' end of SEQ ID NO:151. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:151 from nucleotide 301 to nucleotide 1035, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:151 from nucleotide 301 to nucleotide 1035, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:151 from nucleotide

301 to nucleotide 1035. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:151 from nucleotide 916 to nucleotide 1035, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:151 from nucleotide 916 to nucleotide 1035, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:151 from nucleotide 916 to nucleotide 1035.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:152;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:152, the fragment comprising eight contiguous amino acids of SEQ ID NO:152; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:152. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 20 of SEQ ID NO:152, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:152.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:153;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qf122\_3 deposited under accession number ATCC 207187;
- 30 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qf122\_3 deposited under accession number ATCC 207187;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:154;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:154;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:153.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281; the nucleotide sequence of the full-length protein coding sequence of clone qf122\_3 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone qf122\_3 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:154, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment comprising the amino acid sequence from amino acid 193 to amino acid 202 of SEQ ID NO:154.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:153.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:153; and
- 10 (ab) the nucleotide sequence of the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the
- 15 probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 20 (ba) SEQ ID NO:153; and
- (bb) the nucleotide sequence of the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 25 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:153, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:153 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:153. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281, and extending contiguously from a nucleotide sequence

30

corresponding to the 5' end of said sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:154;
- (b) a fragment of the amino acid sequence of SEQ ID NO:154, the fragment comprising eight contiguous amino acids of SEQ ID NO:154; and
- 10 (c) the amino acid sequence encoded by the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:154. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:154, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment comprising the amino acid sequence from amino acid 193 to amino acid 202 of SEQ ID NO:154.

20 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:155;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742;
- 25 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187;
- 30 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:156;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:156;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:155.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742; the nucleotide sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742; the nucleotide sequence of the full-length protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:156, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:156.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:155.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:155, but excluding the poly(A) tail at the 3' end of SEQ ID NO:155; and

(ab) the nucleotide sequence of the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

20 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:155, but excluding the poly(A) tail at the 3' end of SEQ ID NO:155; and

25 (bb) the nucleotide sequence of the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:155, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:155 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:155, but excluding the poly(A) tail at the 3' end of SEQ ID NO:155. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:156;
- (b) a fragment of the amino acid sequence of SEQ ID NO:156, the fragment comprising eight contiguous amino acids of SEQ ID NO:156; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:156. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 25 of SEQ ID NO:156, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:156.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:157;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757;



- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ys20\_1 deposited under accession number ATCC 207187;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ys20\_1 deposited under accession number ATCC 207187;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:158;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:158;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:157.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757; the nucleotide sequence of the full-length protein coding sequence of clone ys20\_1 deposited under accession number ATCC 207187; or the nucleotide sequence of a mature protein coding sequence of clone ys20\_1 deposited under accession number ATCC 207187. In other preferred embodiments, the
- 30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:158, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158 having biological activity, the fragment comprising the amino acid sequence from amino acid 114 to amino acid 123 of SEQ ID NO:158.

- 5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:157.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:157, but excluding the poly(A) tail at the 3' end of SEQ ID NO:157; and
- 15 (ab) the nucleotide sequence of the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the
- 20 probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from
- 25 the group consisting of:
- (ba) SEQ ID NO:157, but excluding the poly(A) tail at the 3' end of SEQ ID NO:157; and
- (bb) the nucleotide sequence of the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187;
- 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:157, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:157 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:157, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:157. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:158;
- (b) a fragment of the amino acid sequence of SEQ ID NO:158, the fragment comprising eight contiguous amino acids of SEQ ID NO:158; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
ys20\_1 deposited under accession number ATCC 207187;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:158. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:158, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158 having biological activity, the fragment comprising the amino acid sequence from amino acid 114 to amino acid 123 of SEQ ID NO:158.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:159;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:159 from nucleotide 28 to nucleotide 2253;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:159 from nucleotide 568 to nucleotide 2253;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone as180\_1 deposited under accession number ATCC XXXXXX;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone as180\_1 deposited under accession number ATCC XXXXXX;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:160;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:160;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:159.
- 25

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:159 from nucleotide 28 to nucleotide 2253; the nucleotide sequence of SEQ ID NO:159 from nucleotide 568 to nucleotide 2253; the nucleotide sequence of the full-length protein coding sequence of clone as180\_1 deposited under accession number ATCC XXXXXX; or the nucleotide sequence of a mature protein coding sequence of clone as180\_1 deposited under accession number ATCC XXXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX. In further

30

preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:160, or a polynucleotide  
5 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160 having biological activity, the fragment comprising the amino acid sequence from amino acid 366 to amino acid 375 of SEQ ID NO:160.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:159.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
15 consisting of:

(aa) SEQ ID NO:159; and

(ab) the nucleotide sequence of the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;

(ii) hybridizing said probe(s) to human genomic DNA in  
20 conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
25 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:159; and

(bb) the nucleotide sequence of the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;

(ii) hybridizing said primer(s) to human genomic DNA in  
30 conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:159, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:159 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:159. Also

5 preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:159 from nucleotide 28 to nucleotide 2253, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:159 from nucleotide 28 to nucleotide 2253, to a nucleotide sequence corresponding to the 3' end of said sequence of

10 SEQ ID NO:159 from nucleotide 28 to nucleotide 2253. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:159 from nucleotide 568 to nucleotide 2253, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:159 from nucleotide 568 to nucleotide 2253, to a nucleotide

15 sequence corresponding to the 3' end of said sequence of SEQ ID NO:159 from nucleotide 568 to nucleotide 2253.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:160;
- (b) a fragment of the amino acid sequence of SEQ ID NO:160, the fragment comprising eight contiguous amino acids of SEQ ID NO:160; and
- (c) the amino acid sequence encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:160. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 30 of SEQ ID NO:160, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160 having biological activity, the fragment comprising the amino acid sequence from amino acid 366 to amino acid 375 of SEQ ID NO:160.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial,

yeast, insect and mammalian cells, transformed with such polynucleotide compositions. Also provided by the present invention are organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein.

- 5 Processes are also provided for producing a protein, which comprise:
- (a) growing a culture of the host cell transformed with such polynucleotide compositions in a suitable culture medium; and
  - (b) purifying the protein from the culture.

The protein produced according to such methods is also provided by the present  
10 invention.

Protein compositions of the present invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

- 15 Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

- 20 Figures 1A and 1B are schematic representations of the pED6 and pNOTs vectors, respectively, used for deposit of clones disclosed herein.

Figure 2 is a schematic representation of the pCMVSPORT2 vector used for deposit of clone qs14\_3 disclosed herein.

#### DETAILED DESCRIPTION

##### ISOLATED PROTEINS AND POLYNUCLEOTIDES

- 25 Nucleotide and amino acid sequences, as presently determined, are reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone  
30 in accordance with known methods. The predicted amino acid sequence (both full-length and mature forms) can then be determined from such nucleotide sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have

determined to be the reading frame best identifiable with sequence information available at the time of filing.

As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

10        Clone "co62\_12"

A polynucleotide of the present invention has been identified as clone "co62\_12". co62\_12 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. co62\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "co62\_12 protein").

The nucleotide sequence of co62\_12 as presently determined is reported in SEQ ID NO:1, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the co62\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2. Amino acids 1 to 11 of SEQ ID NO:2 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 12. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the co62\_12 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone co62\_12 should be approximately 2200 bp.

The nucleotide sequence disclosed herein for co62\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. co62\_12 demonstrated at least some similarity with sequences identified as AA019597 (ze60f10.s1 Soares retina N2b4HR Homo sapiens cDNA), AA021678 (mh82c02.r1 Soares mouse placenta 4NbMP13.5 14.5 Mus), AA057573 (zf62d10.s1 Soares retina N2b4HR Homo sapiens cDNA clone 381523 3' similar to WP



T12G3.4 CE06440 STRICTOSIDINE SYNTHASE LIKE, mRNA sequence), AA130982, AA287697 (zs53g02.r1 Soares NbHTGBC Homo sapiens cDNA clone 701234 5'), AI042188 (oy37d10.x1 Soares\_parathyroid\_tumor\_NbHPA Homo sapiens cDNA clone IMAGE:1668019 3' similar to WP:F57C2.5 CE16156, mRNA sequence), R63905 (yi19b03.s1  
5 Homo sapiens cDNA clone 139661 3'), T03538 (IB43 Infant brain, Bento Soares Homo sapiens cDNA clone IB43 3'end), T20257 (Human gene signature HUMGS01405), and T23663 (seq294 Homo sapiens cDNA clone b4HB3MA-Cot109+103-Bio-24 3'). The predicted amino acid sequence disclosed herein for co62\_12 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol.  
10 The predicted co62\_12 protein demonstrated at least some similarity to sequences identified as R88502 (Protein sequence for mediating male fertility in plants) and Z83110 (F57C2.5 [Caenorhabditis elegans]). Based upon sequence similarity, co62\_12 proteins and each similar protein or peptide may share at least some activity.

15       Clone "lo311\_8"

A polynucleotide of the present invention has been identified as clone "lo311\_8". lo311\_8 was isolated from a human adult thyroid cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
20 analysis of the amino acid sequence of the encoded protein. lo311\_8 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "lo311\_8 protein").

The nucleotide sequence of lo311\_8 as presently determined is reported in SEQ ID NO:3, and includes a poly(A) tail. What applicants presently believe to be the proper  
25 reading frame and the predicted amino acid sequence of the lo311\_8 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:4. Amino acids 17 to 29 of SEQ ID NO:4 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
30 should the predicted leader/signal sequence not be separated from the remainder of the lo311\_8 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone lo311\_8 should be approximately 3850 bp.

The nucleotide sequence disclosed herein for lo311\_8 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. lo311\_8 demonstrated at least some similarity with sequences identified as AA046836 (zf14b10.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 376891 5' similar to WP:ZK686.3 CE00457), AA297716 (EST113273 Infant adrenal gland, subtracted (total cDNA) I Homo sapiens cDNA 5' end similar to similar to C. elegans hypothetical protein, cosmid ZK686\_3), AF008554 (Rattus norvegicus implantation-associated protein (IAG2) mRNA, partial cds), T68050 (yc39h10.r1 Homo sapiens cDNA clone 83107 5' similar to SP ZK686.3 CE00457), and U42349 (Human N33 mRNA, complete cds). The predicted amino acid sequence disclosed herein for lo311\_8 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted lo311\_8 protein demonstrated at least some similarity to sequences identified as AF008554 (implantation-associated protein [Rattus norvegicus]), R85333 (Human prostate/colon tumour suppressor protein form 1) and U42349 (39 kDa encoded by N33 [Homo sapiens]). Based upon sequence similarity, lo311\_8 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five additional potential transmembrane domains within the lo311\_8 protein sequence, centered around amino acids 10, 190, 220, 275, and 310 of SEQ ID NO:4, respectively.

#### Clone "ns197\_1"

A polynucleotide of the present invention has been identified as clone "ns197\_1". ns197\_1 was isolated from a human adult retina (retinoblastoma WERI-Rb1) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ns197\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ns197\_1 protein").

The nucleotide sequence of ns197\_1 as presently determined is reported in SEQ ID NO:5, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ns197\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:6.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ns197\_1 should be approximately 3650 bp.

The nucleotide sequence disclosed herein for ns197\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ns197\_1 demonstrated at least some similarity with sequences identified as AA495135 (fa03c11.r1 Zebrafish ICRFzfls Danio rerio cDNA clone 3K8 5' similar to WP:ZC518.3 CE06603 ALCOHOL DEHYDROGENASE TRANSCRIPTION EFFECTOR LIKE; mRNA sequence). The predicted amino acid sequence disclosed herein for ns197\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ns197\_1 protein demonstrated at least some similarity to the sequence identified as Z68753 (ZC518.3 [Caenorhabditis elegans]). Based upon sequence similarity, ns197\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the ns197\_1 protein sequence centered around amino acid 135 of SEQ ID NO:6. The nucleotide sequence of ns197\_1 indicates that it may contain one or more repeat sequences, including primate simple repeat GCC, Alu, and other repetitive elements.

#### Clone "pj193\_5"

A polynucleotide of the present invention has been identified as clone "pj193\_5". pj193\_5 was isolated from a human fetal carcinoma (NTD2 cells, treated with retinoic acid for 23 days) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pj193\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pj193\_5 protein").

The nucleotide sequence of pj193\_5 as presently determined is reported in SEQ ID NO:7, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pj193\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:8. Amino acids 9 to 21 of SEQ ID NO:8 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pj193\_5 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pj193\_5 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for pj193\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pj193\_5 demonstrated at least some similarity with sequences identified as AA296889 (EST112653 Cerebellum II Homo sapiens cDNA 5' end), AA296961 (EST112514 Adrenal gland tumor Homo sapiens cDNA 5' end), AA661635 (nv02g06.s1 NCI\_CGAP\_Pr22 Homo sapiens cDNA clone IMAGE:1219066), and U80744 (Homo sapiens CTG4a mRNA, complete cds). The predicted amino acid sequence disclosed herein for pj193\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pj193\_5 protein demonstrated at least some similarity to the sequence identified as U80744 (CTG4a [Homo sapiens]). Based upon sequence similarity, pj193\_5 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of pj193\_5 indicates that it may contain CAG nucleotide repeats; these repeats may create a "hotspot" for certain types of mutations. "Twelve diseases, most with neuropsychiatric features, arise from trinucleotide repeat expansion mutations. Expansion mutations may also cause a number of other disorders, including several additional forms of spinocerebellar ataxia, bipolar affective disorder, schizophrenia, and autism." (Margolis *et al.*, 1997, *Human Genetics* 100(1): 114-122, which is incorporated by reference herein.) It is possible that the gene corresponding to pj193\_5 undergoes a similar etiology.

pj193\_5 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 31 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "pj317\_2"

A polynucleotide of the present invention has been identified as clone "pj317\_2". pj317\_2 was isolated from a human fetal carcinoma (NTD2 cells, treated with retinoic acid for 23 days) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pj317\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pj317\_2 protein").

The nucleotide sequence of pj317\_2 as presently determined is reported in SEQ ID NO:9, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pj317\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:10.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pj317\_2 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for pj317\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pj317\_2 demonstrated at least some similarity with sequences  
10 identified as AA305508 (EST176494 Colon carcinoma (Caco-2) cell line II Homo sapiens cDNA 5' end, mRNA sequence), AA471379 (PMY1151 KG1a Lambda Zap Express cDNA Library Homo sapiens cDNA 5', mRNA sequence), and AA906311 (ok03f08.s1 Soares NFL\_T\_GBC\_S1 Homo sapiens cDNA clone IMAGE:1506759 3', mRNA sequence). The predicted amino acid sequence disclosed herein for pj317\_2 was searched against the  
15 GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pj317\_2 protein demonstrated at least some similarity to the sequences identified as U37763 (Per9p [Pichia angusta]) and U56965 (Caenorhabditis elegans cosmid C15H9). Per9p is a peroxisomal membrane protein, and the predicted pj317\_2 protein demonstrated at least some similarity to peroxisomal proteins from other species as well.  
20 Based upon sequence similarity, pj317\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pj317\_2 protein sequence centered around amino acid 25 of SEQ ID NO:10. The nucleotide sequence of pj317\_2 indicates that it may contain a simple AT and MER repeat region.

25

#### Clone "pt332\_1"

A polynucleotide of the present invention has been identified as clone "pt332\_1". pt332\_1 was isolated from a human adult blood (lymphoblastic leukemia MOLT-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S.  
30 Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pt332\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pt332\_1 protein").

The nucleotide sequence of pt332\_1 as presently determined is reported in SEQ ID NO:11, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pt332\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:12. Amino acids 287 to 299 of SEQ ID NO:12 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 300. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pt332\_1 protein.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pt332\_1 should be approximately 3450 bp.

The nucleotide sequence disclosed herein for pt332\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pt332\_1 demonstrated at least some similarity with sequences identified as AA167221 (zp13c09.s1 Stratagene fetal retina 937202 Homo sapiens cDNA clone 609328 3'), AA437109 (zv53c07.s1 Soares testis NHT Homo sapiens cDNA clone 757356 3'), H14107 (ym62a06.r1 Homo sapiens cDNA clone 163474 5'), and U41264 (C. elegans cDNA). Based upon sequence similarity, pt332\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pt332\_1 protein sequence centered around amino acid 270 of SEQ ID NO:12.

20 pt332\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 100 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

25

#### Clone "qc297\_15"

A polynucleotide of the present invention has been identified as clone "qc297\_15". qc297\_15 was isolated from a human adult neural (neuroepithelioma HTB-10 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qc297\_15 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qc297\_15 protein").

The nucleotide sequence of qc297\_15 as presently determined is reported in SEQ ID NO:13, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qc297\_15 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:14.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qc297\_15 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for qc297\_15 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qc297\_15 demonstrated at least some similarity with sequences  
10 identified as AA625537 (af72g07.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 1047612 5') and T24537 (EST112 Homo sapiens cDNA clone 4H3). Based upon sequence similarity, qc297\_15 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the qc297\_15 protein sequence, around amino acid 20 of SEQ ID NO:14. The  
15 nucleotide/amino acid sequence of qc297\_15 indicates that it may contain an Alu/SVA/MER repeat region.

qc297\_15 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 7 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

20

#### Clone "qg596\_12"

A polynucleotide of the present invention has been identified as clone "qg596\_12". qg596\_12 was isolated from a human adult neural (neuroepithelioma HTB-10 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins  
25 (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qg596\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qg596\_12 protein").

The nucleotide sequence of qg596\_12 as presently determined is reported in SEQ  
30 ID NO:15, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qg596\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:16.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qg596\_12 should be approximately 2750 bp.

The nucleotide sequence disclosed herein for qg596\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qg596\_12 demonstrated at least some similarity with sequences identified as AA332939 (EST37132 Embryo, 8 week I Homo sapiens cDNA 5' end),  
5 AA334678 (EST39190 Embryo, 9 week Homo sapiens cDNA 5' end), AA362653 (EST72375 Namalwa B cells I Homo sapiens cDNA 5' end), and AA829841 (od40d01.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1370401 3' similar to WP:F10G7.1 CE02624). The predicted amino acid sequence disclosed herein for qg596\_12 was searched  
10 against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qg596\_12 protein demonstrated at least some similarity to the sequence identified as U40029 (coded for by C. elegans cDNA yk16b1.3; coded for by C. elegans cDNA yk8g6.5; coded for by C. elegans cDNA yk8g6.3; coded for by C. elegans cDNA yk6d3.5). Based upon sequence similarity, qg596\_12 proteins and each  
15 similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the qg596\_12 protein sequence, one centered around amino acid 180 and another around amino acid 660 of SEQ ID NO:16.

qg596\_12 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 33 kDa was detected in membrane fractions using SDS  
20 polyacrylamide gel electrophoresis.

#### Clone "rb649\_3"

A polynucleotide of the present invention has been identified as clone "rb649\_3". rb649\_3 was isolated from a human fetal kidney (293 cell line) cDNA library using  
25 methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rb649\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rb649\_3 protein").

30 The nucleotide sequence of rb649\_3 as presently determined is reported in SEQ ID NO:17, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rb649\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:18. Amino acids 42 to 54 of SEQ ID NO:18 are a predicted leader/signal sequence, with the predicted



mature amino acid sequence beginning at amino acid 55. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the rb649\_3 protein.

- 5       The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rb649\_3 should be approximately 2500 bp.

The nucleotide sequence disclosed herein for rb649\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rb649\_3 demonstrated at least some similarity with sequences  
10 identified as AA177001 (nc01h02.s1 NCI\_CGAP\_Pr1 Homo sapiens cDNA clone IMAGE 182). The predicted amino acid sequence disclosed herein for rb649\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted rb649\_3 protein demonstrated at least some similarity to  
15 sequences identified as AB002405 (LAK-4p [Homo sapiens]), R89470 (Collagen/TGF-beta-1 fusion protein), and U23516 (Undefined [Caenorhabditis elegans]). Based upon sequence similarity, rb649\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts eight additional potential transmembrane domains within the rb649\_3 protein sequence, centered around amino acids 140, 240, 280, 325, 370, 425, 475, and 540 of SEQ ID NO:18, respectively. The  
20 nucleotide sequence of rb649\_3 indicates that it may contain a simple GGA repeat region.

#### Clone "ca106\_19x"

A polynucleotide of the present invention has been identified as clone "ca106\_19x". A cDNA clone was first isolated from a mouse embryonic (ES cell embryoid bodies  
25 harvested 2-12 days after LIF removed) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate ca106\_19x from a mixture of human fetal brain and human adult brain cDNA libraries.  
30 ca106\_19x is a full-length human clone, including the entire coding sequence of a secreted protein (also referred to herein as "ca106\_19x protein").

The nucleotide sequence of ca106\_19x as presently determined is reported in SEQ ID NO:19, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the ca106\_19x protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:20.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ca106\_19x should be approximately 4050 bp.

5       The nucleotide sequence disclosed herein for ca106\_19x was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ca106\_19x demonstrated at least some similarity with sequences identified as AA886998 (oj30g03.s1 NCI\_CGAP\_Lu5 Homo sapiens cDNA clone IMAGE:1499860 3'), F08279 (H. sapiens partial cDNA sequence; clone c-zpe11), F13022 (H.  
10 sapiens partial cDNA sequence; clone c-3hf07), H38128 (yp46c12.s1 Homo sapiens cDNA clone 190486 3'), T77601 (yc91e07.r1 Homo sapiens cDNA clone 23192 5'), U93720 (Homo sapiens TEX28 mRNA, complete cds), W55512 (ma28h03.r1 Life Tech mouse brain Mus musculus cDNA clone 312053 5'), and Z22333 (H.sapiens DNA sequence). The predicted amino acid sequence disclosed herein for ca106\_19x was searched against the GenPept  
15 and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ca106\_19x protein demonstrated at least some similarity to sequences identified as U56965 (C15H9.4 gene product [Caenorhabditis elegans]) and U93720 (TEX28 [Homo sapiens]). Based upon sequence similarity, ca106\_19x proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts  
20 four potential transmembrane domains within the ca106\_19x protein sequence, centered around amino acids 170, 430, 590, and 625 of SEQ ID NO:20, respectively. The nucleotide sequence of ca106\_19x indicates that it contains at least one repetitive element.

#### Clone "ci52\_2"

25       A polynucleotide of the present invention has been identified as clone "ci52\_2". A cDNA clone was first isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then  
30 used to isolate ci52\_2 from a human fetal brain cDNA library. ci52\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ci52\_2 protein").

The nucleotide sequence of ci52\_2 as presently determined is reported in SEQ ID NO:21, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the ci52\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:22. Amino acids 9 to 21 of SEQ ID NO:22 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ci52\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ci52\_2 should be approximately 1775 bp.

10 The nucleotide sequence disclosed herein for ci52\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ci52\_2 demonstrated at least some similarity with sequences identified as AA083339 (zn31d10.r1 Stratagene endothelial cell 937223 Homo sapiens cDNA clone 549043 5'), AA514339 (nf56c10.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA  
15 clone 923922), AA628942 (af28f01.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 1032985 3', mRNA sequence), M78692 (EST00840 Homo sapiens cDNA clone HHCMC16), N67265 (yz49d04.s1 Homo sapiens cDNA clone 286375 3'), N95514 (yy62d10.r1 Homo sapiens cDNA clone 278131 5'), Q60715 (Human brain Expressed Sequence Tag EST00840; standard; cDNA), and R46588 (yg51a12.s1 Homo sapiens cDNA  
20 clone 35984 3'). The predicted amino acid sequence disclosed herein for ci52\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ci52\_2 protein demonstrated at least some similarity to the sequence identified as M68866 (stranded at second [Drosophila melanogaster]). Based upon sequence similarity, ci52\_2 proteins and each similar protein  
25 or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the ci52\_2 protein sequence, one around amino acid 146 and another around amino acid 177 of SEQ ID NO:22.

#### Clone "md124\_16"

30 A polynucleotide of the present invention has been identified as clone "md124\_16". A cDNA clone was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then

used to isolate md124\_16 from a human adult kidney cDNA library. md124\_16 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "md124\_16 protein").

5 The nucleotide sequence of md124\_16 as presently determined is reported in SEQ ID NO:23, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the md124\_16 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:24. Amino acids 152 to 164 of SEQ ID NO:24 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 165. Due to the  
10 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the md124\_16 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone md124\_16 should be approximately 2300 bp.

15 The nucleotide sequence disclosed herein for md124\_16 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. md124\_16 demonstrated at least some similarity with sequences identified as AA215643 (zr98d05.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:683721 3'), AA489121 (aa56b07.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone  
20 IMAGE:824917 5'), W72865 (zd59e07.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344964 3'), and W76100 (zd59e07.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344964 5'). Based upon sequence similarity, md124\_16 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of md124\_16 indicates that it may contain at least one MER repeat sequence.  
25

#### Clone "pk366\_7"

A polynucleotide of the present invention has been identified as clone "pk366\_7". pk366\_7 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No.  
30 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pk366\_7 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pk366\_7 protein").

The nucleotide sequence of pk366\_7 as presently determined is reported in SEQ ID NO:25, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pk366\_7 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:26. Amino acids 361 to 373 of SEQ ID NO:26 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 374. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pk366\_7 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pk366\_7 should be approximately 3300 bp.

The nucleotide sequence disclosed herein for pk366\_7 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pk366\_7 demonstrated at least some similarity with sequences identified as AA057428 (zf57c11.s1 Soares retina N2b4HR Homo sapiens cDNA clone 381044 3'), AA457625 (aa89e09.r1 Stratagene fetal retina 937202 Homo sapiens cDNA clone 838504 5'), AA601545 (nn87h11.s1 NCI\_CGAP\_Br2 Homo sapiens cDNA clone IMAGE:1098213), T19564 (Human gene signature HUMGS00629; standard; cDNA to mRNA), and U94831 (Homo sapiens multispinning membrane protein mRNA, complete cds). The predicted amino acid sequence disclosed herein for pk366\_7 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pk366\_7 protein demonstrated at least some similarity to sequences identified as D87444 (endomembrane protein EMP70 precursor isolog [Arabidopsis thaliana]), U94831 (multispinning membrane protein [Homo sapiens]), and U95973 (endomembrane protein EMP70 precursor isolog [Arabidopsis thaliana]). Based upon sequence similarity, pk366\_7 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts nine additional potential transmembrane domains within the pk366\_7 protein sequence, centered around amino acids 191, 260, 288, 325, 355, 412, 447, 481, and 517 of SEQ ID NO:26, respectively.

#### Clone "pl741\_5"

A polynucleotide of the present invention has been identified as clone "pl741\_5". pl741\_5 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No.

5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pl741\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pl741\_5 protein").

5 The nucleotide sequence of pl741\_5 as presently determined is reported in SEQ ID NO:27, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pl741\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28. Amino acids 3 to 15 of SEQ ID NO:28 are a predicted leader/signal sequence, with the predicted  
10 mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pl741\_5 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
15 pl741\_5 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for pl741\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pl741\_5 demonstrated at least some similarity with sequences identified as AA283176 (zt17a04.s1 Soares ovary tumor NbHOT Homo sapiens cDNA  
20 clone 713358 3'), AA204801 (zq61d12.r1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 646103 5'), and H59410 (yr19g04.r1 Homo sapiens cDNA clone 205782 5'). The predicted amino acid sequence disclosed herein for pl741\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pl741\_5 protein demonstrated at least some similarity to  
25 sequences identified as U00027 (Cdc23p cell cycle protein [Saccharomyces cerevisiae]) and U58763 (F10C5.1 [Caenorhabditis elegans]). Based upon sequence similarity, pl741\_5 proteins and each similar protein or peptide may share at least some activity. Analysis of the amino acid sequence of the predicted pl741\_5 protein reveals the presence of four TPR (tetratricopeptide) domains. TPR domains are found in a wide variety of proteins with  
30 varying functions and localizations — from the nucleus to the extracellular milieu — and are thought to function as protein-protein interaction domains. The TPR domains are found at amino acid residues 166-194, 328-356, 362-390, and 396-424 of SEQ ID NO:28.

Clone "pp314\_19"

A polynucleotide of the present invention has been identified as clone "pp314\_19". pp314\_19 was isolated from a human adult blood (lymphoblastic leukemia MOLT-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pp314\_19 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pp314\_19 protein").

The nucleotide sequence of pp314\_19 as presently determined is reported in SEQ ID NO:29, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pp314\_19 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:30. Amino acids 147 to 159 of SEQ ID NO:30 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 160; amino acids 238 to 250 of SEQ ID NO:30 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 251. Due to the hydrophobic nature of these possible leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the pp314\_19 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pp314\_19 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for pp314\_19 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pp314\_19 demonstrated at least some similarity with sequences identified as AA044042 (zk58g05.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 487064 5', mRNA sequence), AA127902 (zl12d01.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 501697 5'), AA609481 (af14a12.s1 Soares testis NHT Homo sapiens cDNA clone 1031614 3', mRNA sequence), T26699 (Human gene signature HUMGS08949; standard; cDNA to mRNA), and W93399 (zd95b06.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 357203 3'). The predicted amino acid sequence disclosed herein for pp314\_19 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pp314\_19 protein demonstrated at least some similarity to sequences identified as AE000857 (chaperonin [Methanobacterium thermoautotrophicum]), AJ006549 (ThsA [Pyrodictium occultum]), and L34691 (thermophilic factor 56 [Sulfolobus shibatae]). Based upon sequence

similarity, pp314\_19 proteins and each similar protein or peptide may share at least some activity. Analysis of the amino acid sequence of the predicted pp314\_19 protein revealed the cpn60\_TCP1 signature (at amino acids 29-570 of SEQ ID NO:30) which has some ATPase activity and is indicative of chaperonins. A P-loop motif — a common motif in ATP- and GTP-binding proteins — is found around amino acid 200 of SEQ ID NO:30. The presence of the P-loop is interesting when taken in conjunction with the potential ATPase activity associated with the cpn60\_TCP1 signature. The TopPredII computer program predicts three additional potential transmembrane domains within the pp314\_19 protein sequence, centered around amino acids 55, 90, and 330 of SEQ ID NO:30, respectively.

pp314\_19 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 6 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

15      Clone "pv35\_1"

A polynucleotide of the present invention has been identified as clone "pv35\_1". pv35\_1 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pv35\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pv35\_1 protein").

The nucleotide sequence of pv35\_1 as presently determined is reported in SEQ ID NO:31, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pv35\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:32.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pv35\_1 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for pv35\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pv35\_1 demonstrated at least some similarity with sequences identified as AA335869 (EST40348 Epididymus Homo sapiens cDNA 5' end), AA599418 (ag23c03.s1 Jia bone marrow stroma Homo sapiens cDNA clone 1071172 3'), and H03595



(yj42e06.r1 Homo sapiens cDNA clone 151426 5'). The predicted amino acid sequence disclosed herein for pv35\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pv35\_1 protein demonstrated at least some similarity to sequences identified as Z99277 (Y53C12A.3 [Caenorhabditis elegans]). Based upon sequence similarity, pv35\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four potential transmembrane domains within the pv35\_1 protein sequence, centered around amino acids 127, 161, 192, and 250 of SEQ ID NO:32, respectively.

10

#### Clone "pw337\_6"

A polynucleotide of the present invention has been identified as clone "pw337\_6". pw337\_6 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pw337\_6 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pw337\_6 protein").

The nucleotide sequence of pw337\_6 as presently determined is reported in SEQ ID NO:33, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pw337\_6 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:34. Another potential pw337\_6 reading frame and predicted amino acid sequence is encoded by basepairs 648 to 908 of SEQ ID NO:33 and is reported in SEQ ID NO:238. The overlapping reading frames of SEQ ID NO:34 and SEQ ID NO:238 could be joined if a frameshift were introduced into the nucleotide sequence of SEQ ID NO:33 between position 645 and position 736.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pw337\_6 should be approximately 1000 bp.

The nucleotide sequence disclosed herein for pw337\_6 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pw337\_6 demonstrated at least some similarity with sequences identified as AA682471 (zj18c02.s1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 450626 3', mRNA sequence), T20708 (Human gene signature HUMGS01925;

standard; cDNA to mRNA), W24658 (zb63b05.r1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 308241 5'), and Z82192 (Homo sapiens DNA sequence from PAC 186O1 on chromosome 22). The predicted amino acid sequence disclosed herein for pw337\_6 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pw337\_6 protein demonstrated at least some similarity to the sequence identified as Z82192 (dJ186O1.1 [Homo sapiens]). Based upon sequence similarity, pw337\_6 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pw337\_6 protein sequence centered around amino acid 75 of SEQ ID NO:34. The nucleotide sequence of pw337\_6 indicates that it may contain one or more repetitive elements.

pw337\_6 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 22 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "rd610\_1"

A polynucleotide of the present invention has been identified as clone "rd610\_1". rd610\_1 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rd610\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rd610\_1 protein").

The nucleotide sequence of rd610\_1 as presently determined is reported in SEQ ID NO:35, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rd610\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:36.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rd610\_1 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for rd610\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rd610\_1 demonstrated at least some similarity with sequences identified as AA442056 (zw56f08.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 774087 5'), AA992905 (ot92b06.s1 Soares\_total\_fetus\_Nb2HF8\_9w Homo sapiens

cDNA clone IMAGE 1624211 3', mRNA sequence), D31767 (Human mRNA for KIAA0058 gene, complete cds), and T40090 (Human Serrate-1 (HJ1) cDNA; standard; cDNA). Based upon sequence similarity, rd610\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential  
5 transmembrane domain within the rd610\_1 protein sequence centered around amino acid 30 of SEQ ID NO:36; amino acids 23 to 35 of SEQ ID NO:36 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 36.

rd610\_1 protein was expressed in a COS cell expression system, and an expressed  
10 protein band of approximately 7 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

#### Clone "rd810\_6"

A polynucleotide of the present invention has been identified as clone "rd810\_6".  
15 rd810\_6 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rd810\_6 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to  
20 herein as "rd810\_6 protein").

The nucleotide sequence of rd810\_6 as presently determined is reported in SEQ ID NO:37, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rd810\_6 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:38. Amino  
25 acids 112 to 124 of SEQ ID NO:38 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 125. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the rd810\_6 protein.

30 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rd810\_6 should be approximately 850 bp.

The nucleotide sequence disclosed herein for rd810\_6 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rd810\_6 demonstrated at least some similarity with sequences

identified as AA452718 (zx39d04.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 788839 5', mRNA sequence), AA292888 (zt66c01.r1 Soares testis NHT Homo sapiens cDNA clone 727296 5'), and T23348 (Human gene signature HUMGS05169; standard; cDNA to mRNA). Based upon sequence similarity, rd810\_6 proteins and each similar  
5 protein or peptide may share at least some activity.

rd810\_6 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 23 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### 10 Clone "cf85\_1"

A polynucleotide of the present invention has been identified as clone "cf85\_1". A cDNA clone was first isolated from a human adult placenta library cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis  
15 of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate cf85\_1 from a human adult brain cDNA library. cf85\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cf85\_1 protein").

The nucleotide sequence of cf85\_1 as presently determined is reported in SEQ ID NO:39, and includes a poly(A) tail. What applicants presently believe to be the proper  
20 reading frame and the predicted amino acid sequence of the cf85\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:40.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cf85\_1 should be approximately 2000 bp.

25 The nucleotide sequence disclosed herein for cf85\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cf85\_1 demonstrated at least some similarity with sequences identified as H50932 (yo35f03.r1 Homo sapiens cDNA clone 179933 5'), H51595 (yo35f03.s1 Homo sapiens cDNA clone 179933 3'), and T24664 (Human gene signature HUMGS06728; standard; cDNA to mRNA). Based upon sequence similarity, cf85\_1  
30 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three potential transmembrane domains within the cf85\_1 protein sequence, centered around amino acids 150, 195, and 220 of SEQ ID NO:40,

respectively. The nucleotide sequence of cf85\_1 indicates that it may contain an Alu repetitive element.

Clone "dd504\_18"

5 A polynucleotide of the present invention has been identified as clone "dd504\_18". dd504\_18 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dd504\_18 is a full-length  
10 clone, including the entire coding sequence of a secreted protein (also referred to herein as "dd504\_18 protein").

The nucleotide sequence of dd504\_18 as presently determined is reported in SEQ ID NO:41, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dd504\_18 protein  
15 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:42. Amino acids 134 to 146 of SEQ ID NO:42 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 147. Amino acids 7 to 19 of SEQ ID NO:42 are also a possible leader/signal sequence, with a predicted mature amino acid sequence beginning in that case at amino acid 20. Due to the hydrophobic  
20 nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the dd504\_18 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dd504\_18 should be approximately 2000 bp.

The nucleotide sequence disclosed herein for dd504\_18 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dd504\_18 demonstrated at least some similarity with sequences identified as AA393779 (zt77f07.r1 Soares testis NHT Homo sapiens cDNA clone 728389 5' similar to WP:F41E7.1 CE03301; mRNA sequence), AA429420 (zw51f02.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773595 5' similar to WP W02B12.7 CE03767  
30 KINENSIN-LIKE PROTEIN), AC002038 (\*\* SEQUENCING IN PROGRESS \*\*) Human chromosome 16p12 BAC clone CIT987SK-101B6; HTGS phase 1, 1 unordered pieces; Homo sapiens chromosome 2 clone 101B6 from 2p11, complete sequence), H10672 (yl99g09.r1 Homo sapiens cDNA clone 46448 5'), and R59895 (yh07f12.r1 Homo sapiens cDNA clone 42477 5'). The predicted amino acid sequence disclosed herein for dd504\_18

was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dd504\_18 protein demonstrated at least some similarity to sequences identified as AE000854 (Na<sup>+</sup>/H<sup>+</sup>-exchanging protein Na<sup>+</sup>/H<sup>+</sup> antiporter [*Methanobacterium thermoautotrophicum*]) and Z68106 (F41E7.1 [*Caenorhabditis elegans*]). Based upon sequence similarity, dd504\_18 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts eight potential transmembrane domains within the dd504\_18 protein sequence, centered around amino acids 20, 48, 118, 144, 191, 220, 268, and 326 of SEQ ID NO:42, respectively.

dd504\_18 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 36 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "np26\_3"

A polynucleotide of the present invention has been identified as clone "np26\_3". np26\_3 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. np26\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "np26\_3 protein").

The nucleotide sequence of np26\_3 as presently determined is reported in SEQ ID NO:43, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the np26\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:44.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone np26\_3 should be approximately 3800 bp.

The nucleotide sequence disclosed herein for np26\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. np26\_3 demonstrated at least some similarity with sequences identified as AA118527 (mo99d08.r1 Stratagene mouse heart (#937316) *Mus musculus* cDNA clone 567855 5'), AA284633 (zt15d04.s1 NCI\_CGAP\_GCB1 *Homo sapiens* cDNA clone IMAGE:713191 3', mRNA sequence), AA427620 (zw30d02.s1 Soares ovary tumor NbHOT *Homo sapiens* cDNA clone 770787 3' similar to contains MER17.b1 MER17

repetitive element; mRNA sequence), and AA496955 (aa42f01.s1 Soares NhHMPu S1 Homo sapiens cDNA clone 823609 3', mRNA sequence). The predicted amino acid sequence disclosed herein for np26\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted np26\_3 protein demonstrated at least some similarity to the sequence identified as M86752 (transformation-sensitive protein [Homo sapiens]). Based upon sequence similarity, np26\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the np26\_3 protein sequence centered around amino acid 146 of SEQ ID NO:44.

np26\_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 63 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

#### Clone "pm412\_12"

A polynucleotide of the present invention has been identified as clone "pm412\_12". pm412\_12 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pm412\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm412\_12 protein").

The nucleotide sequence of pm412\_12 as presently determined is reported in SEQ ID NO:45, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pm412\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:46. Amino acids 607 to 619 of SEQ ID NO:46 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 620. Due to the hydrophobic nature of this possible leader/signal sequence, it is likely to act as a transmembrane domain should it not be separated from the remainder of the pm412\_12 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm412\_12 should be approximately 4000 bp.

The nucleotide sequence disclosed herein for pm412\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and

FASTA search protocols. pm412\_12 demonstrated at least some similarity with sequences identified as AA176820 (zp34a12.s1 Stratagene muscle 937209 Homo sapiens cDNA clone 611326 3'), AA425762 (zw47f10.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773227 3' similar to TR:G285999 G285999 ORF, COMPLETE CDS), AA568580 (nm21a10.s1 NCI\_CGAP\_Co10 Homo sapiens cDNA clone IMAGE:1060794 similar to TR:G642306 G642306 HYPOTHETICAL 153.8 KD PROTEIN), AA610863 (np98h01.s1 NCI\_CGAP\_Lu1 Homo sapiens cDNA clone IMAGE 1142449 similar to TR G285999 G285999 ORF, COMPLETE CDS), AA769312 (nz39f06.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 1290179 similar to TR Q15393 Q15393 ORF, COMPLETE CDS; mRNA sequence), D13642 (Human mRNA for KIAA0017 gene, complete cds), and T92977 (ye22e09.r1 Homo sapiens cDNA clone 118504 5'). The predicted amino acid sequence disclosed herein for pm412\_12 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pm412\_12 protein demonstrated at least some similarity to sequences identified as AF043699 (ORF; similar to human UV-damaged DNA binding factor [*C. elegans*]), D13642 (KIAA0017 [*Homo sapiens*]), R72386 (XAP-1, part of the DNA repair complex), and X54413 (alpha1(IX) collagen precursor [*Homo sapiens*]). Based upon sequence similarity, pm412\_12 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three potential transmembrane domains within the pm412\_12 protein sequence, centered around amino acids 277, 415, and 1060 of SEQ ID NO46, respectively.

pm412\_12 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 119 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "pm421\_3"

A polynucleotide of the present invention has been identified as clone "pm421\_3". pm421\_3 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pm421\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm421\_3 protein").



The nucleotide sequence of pm421\_3 as presently determined is reported in SEQ ID NO:47, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pm421\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Amino acids 10 to 22 of SEQ ID NO:48 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pm421\_3 protein.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm421\_3 should be approximately 2800 bp.

The nucleotide sequence disclosed herein for pm421\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pm421\_3 demonstrated at least some similarity with sequences identified as AA196485 (zq59a06.s1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 645874 3'), AA421712 (zu26g11.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 739172 5', mRNA sequence), AC005026 (Homo sapiens clone GS489L14; HTGS phase 1, 3 unordered pieces), AC005028 (Homo sapiens clone GS539F22; HTGS phase 1, 1 unordered pieces), Q60534 (Human brain Expressed Sequence Tag EST02540; standard; cDNA), and R13985 (yf68h04.r1 Homo sapiens cDNA clone 27722 5'). Based upon sequence similarity, pm421\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the pm421\_3 protein sequence centered around amino acid 36 of SEQ ID NO:48.

25

#### Clone "pv6\_1"

A polynucleotide of the present invention has been identified as clone "pv6\_1". pv6\_1 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pv6\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pv6\_1 protein").

30

The nucleotide sequence of pv6\_1 as presently determined is reported in SEQ ID NO:49, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pv6\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:50. Amino acids 39 to 51 of SEQ ID NO:50 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 52. Amino acids 8 to 20 of SEQ ID NO:50 are also a possible leader/signal sequence, with a predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the pv6\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pv6\_1 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for pv6\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pv6\_1 demonstrated at least some similarity with sequences identified as B53192 (CIT-HSP-2009D9.TR CIT-HSP Homo sapiens genomic clone 2009D9, genomic survey sequence), R18429 (yg02g05.r1 Homo sapiens cDNA clone 31056 5'), T77089 (yc93b02.r1 Homo sapiens cDNA clone 23653 5'), and X89480 (S.scrofa mRNA for membrane protein). The predicted amino acid sequence disclosed herein for pv6\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pv6\_1 protein demonstrated at least some similarity to the sequence identified as X89480 (transmembrane protein [Sus scrofa]). Based upon sequence similarity, pv6\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pv6\_1 protein sequence centered around amino acid 21 of SEQ ID NO:50.

#### Clone "qs14\_3"

A polynucleotide of the present invention has been identified as clone "qs14\_3". A cDNA clone was isolated from a human whole embryo cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate qs14\_3 from a human fetal heart cDNA library. qs14\_3 is a full-length

clone, including the entire coding sequence of a secreted protein (also referred to herein as "qs14\_3 protein").

The nucleotide sequence of qs14\_3 as presently determined is reported in SEQ ID NO:51, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the qs14\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:52. Amino acids 15 to 27 of SEQ ID NO:52 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
10 the predicted leader/signal sequence not be separated from the remainder of the qs14\_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qs14\_3 should be approximately 5000 bp.

The nucleotide sequence disclosed herein for qs14\_3 was searched against the  
15 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qs14\_3 demonstrated at least some similarity with sequences identified as AA558554 (nl69g02.s1 NCI\_CGAP\_Pr4.1 Homo sapiens cDNA clone IMAGE 1045970 similar to TR G307329 G307329 PROTOCADHERIN 43), AB002343 (Human mRNA for KIAA0345 gene), and L43592 (Rattus norvegicus protocadherin-3 (pcdh3)  
20 mRNA, and translated products). The predicted amino acid sequence disclosed herein for qs14\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qs14\_3 protein demonstrated at least some similarity to sequences identified as AF029343 (protocadherin [Homo sapiens]), AF042192 (protocadherin [Xenopus]), AF052685 (protocadherin 43 [Homo sapiens]),  
25 L11373 (protocadherin 43 [Homo sapiens]), R49144 (Product of alternative splicing of human protocadherin-43 mRNA), and Y08715 (protocadherin [Mus musculus]). The cadherins are a family of calcium-binding membrane glycoproteins. Most cadherins are capable of acting as cell adhesion molecules (CAMs). Motif analysis of the predicted qs14\_3 protein also detects the 'cadherins extracellular repeated domain signature'. Based  
30 upon sequence similarity, qs14\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the qs14\_3 protein sequence, one centered around amino acid 510 and another around amino acid 721 of SEQ ID NO:52.

qs14\_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 132 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5        Clone "qy338\_9"

A polynucleotide of the present invention has been identified as clone "qy338\_9". qy338\_9 was isolated from a human adult blood (promyelocytic leukemia HL-60) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qy338\_9 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qy338\_9 protein").

The nucleotide sequence of qy338\_9 as presently determined is reported in SEQ ID NO:53, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qy338\_9 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:54. Amino acids 144 to 156 of SEQ ID NO:54 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 157. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qy338\_9 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qy338\_9 should be approximately 1300 bp.

The nucleotide sequence disclosed herein for qy338\_9 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qy338\_9 demonstrated at least some similarity with sequences identified as AA205412 (zq66a09.s1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 646552 3' similar to contains Alu repetitive element; contains element LTR1 repetitive element; mRNA), AA595068 (no40h10.s1 NCI\_CGAP\_Pr23 Homo sapiens cDNA clone IMAGE 1103203 similar to WP C27F2.4 CE01171 METHYLTRANSFERASE), AJ224442 (Homo sapiens mRNA for putative methyltransferase), and H40834 (yo05g09.r1 Homo sapiens cDNA clone 177088 5'). The predicted amino acid sequence disclosed herein for qy338\_9 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qy338\_9 protein

demonstrated at least some similarity to sequences identified as AJ224442 (methyltransferase [Homo sapiens]), U40419 (similar to *S. cerevisiae* gene YCR47C, putative 30.7 kd methyltransferase (SP YCT7\_YEAST,P25627) [*Caenorhabditis elegans*]), and Z69240 (putative methyltransferase [*S. cerevisiae*]). Based upon sequence similarity, qy338\_9  
5 proteins and each similar protein or peptide may share at least some activity.

qy338\_9 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 34 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

10 Clone "rc58\_1"

A polynucleotide of the present invention has been identified as clone "rc58\_1". rc58\_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
15 analysis of the amino acid sequence of the encoded protein. rc58\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rc58\_1 protein").

The nucleotide sequence of rc58\_1 as presently determined is reported in SEQ ID NO:55, and includes a poly(A) tail. What applicants presently believe to be the proper  
20 reading frame and the predicted amino acid sequence of the rc58\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56. Amino acids 2 to 14 of SEQ ID NO:56 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
25 the predicted leader/signal sequence not be separated from the remainder of the rc58\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rc58\_1 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for rc58\_1 was searched against the  
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rc58\_1 demonstrated at least some similarity with sequences identified as AA203670 (zx52d04.r1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 446119 5' similar to gb X07868\_rna1 PUTATIVE INSULIN-LIKE GROWTH FACTOR II ASSOCIATED (HUMAN); mRNA sequence), AA878778 (oe80h01.s1

NCI\_CGAP\_Lu5 Homo sapiens cDNA clone IMAGE:1417969 3', mRNA sequence), and U96448 (Bos taurus cleavage and polyadenylation specificity factor 30 kDa subunit mRNA, complete cds). The predicted amino acid sequence disclosed herein for rc58\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted rc58\_1 protein demonstrated at least some similarity to sequences identified as AF033201 (cleavage and polyadenylation specificity factor [Mus musculus]) and U96448 (cleavage and polyadenylation specificity factor 30 kDa subunit [Bos taurus]). Based upon sequence similarity, rc58\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the rc58\_1 protein sequence centered around amino acid 53 of SEQ ID NO:56.

#### Clone "rd232\_5"

A polynucleotide of the present invention has been identified as clone "rd232\_5". rd232\_5 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rd232\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rd232\_5 protein").

The nucleotide sequence of rd232\_5 as presently determined is reported in SEQ ID NO:57, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rd232\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:58.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rd232\_5 should be approximately 3800 bp.

The nucleotide sequence disclosed herein for rd232\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rd232\_5 demonstrated at least some similarity with sequences identified as AA768103 (oc16g01.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1341072), AA831487 (oc61a11.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1354172 3', mRNA sequence), and R57296 (F2616 Fetal heart Homo sapiens cDNA clone F2616 5' end). The predicted amino acid sequence disclosed herein for rd232\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the

BLASTX search protocol. The predicted rd232\_5 protein demonstrated at least some similarity to the sequence identified as Z79755 (F43G9.2 [Caenorhabditis elegans]). Based upon sequence similarity, rd232\_5 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential  
5 transmembrane domain within the rd232\_5 protein sequence centered around amino acid 225 of SEQ ID NO:58. The nucleotide sequence of rd232\_5 indicates that it may contain a simple AC repeat region.

#### Clone "ck213\_12"

10 A polynucleotide of the present invention has been identified as clone "ck213\_12". ck213\_12 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ck213\_12 is a full-length  
15 clone, including the entire coding sequence of a secreted protein (also referred to herein as "ck213\_12 protein").

The nucleotide sequence of ck213\_12 as presently determined is reported in SEQ ID NO:59, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ck213\_12 protein  
20 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:60.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ck213\_12 should be approximately 3500 bp.

The nucleotide sequence disclosed herein for ck213\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
25 FASTA search protocols. ck213\_12 demonstrated at least some similarity with sequences identified as AA062731 (zm01h03.s1 Stratagene corneal stroma (#937222) Homo sapiens cDNA clone 512885 3' similar to TR:G1136390 G1136390 KIAA0164 PROTEIN, mRNA sequence), AA173803 (zp30f05.s1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 610977 3', mRNA sequence), D79986 (Human mRNA for KIAA0164 protein  
30 gene, complete cds), and R01411 (ye77c11.s1 Homo sapiens cDNA clone 123764 3'). The predicted amino acid sequence disclosed herein for ck213\_12 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ck213\_12 protein demonstrated at least some similarity to the sequence identified as D79986 (similar to human DNA-binding protein 5 [Homo sapiens],

KIAA0164 protein [Homo sapiens], HUMKIAA04\_1). Based upon sequence similarity, ck213\_12 proteins and each similar protein or peptide may share at least some activity.

Clone "pg195\_1"

5 A polynucleotide of the present invention has been identified as clone "pg195\_1". pg195\_1 was isolated from a human adult thyroid cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pg195\_1 is a full-length clone,  
10 including the entire coding sequence of a secreted protein (also referred to herein as "pg195\_1 protein").

The nucleotide sequence of pg195\_1 as presently determined is reported in SEQ ID NO:61, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pg195\_1 protein  
15 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:62.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pg195\_1 should be approximately 3300 bp.

The nucleotide sequence disclosed herein for pg195\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
20 FASTA search protocols. pg195\_1 demonstrated at least some similarity with sequences identified as H72617 (yu02g10.r1 Homo sapiens cDNA clone 232674 5') and W37280 (zc11a07.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 321972 5', mRNA sequence). The predicted amino acid sequence disclosed herein for pg195\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
25 BLASTX search protocol. The predicted pg195\_1 protein demonstrated at least some similarity to the sequence identified as AF007270 (contains similarity to myosin heavy chain [Arabidopsis thaliana]). Based upon sequence similarity, pg195\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the pg195\_1 protein  
30 sequence, one centered around amino acid 480 and another around amino acid 520 of SEQ ID NO:62. The nucleotide sequence of pg195\_1 indicates that it may contain one or more repetitive sequences.



Clone "pw460\_5"

A polynucleotide of the present invention has been identified as clone "pw460\_5". pw460\_5 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pw460\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pw460\_5 protein").

The nucleotide sequence of pw460\_5 as presently determined is reported in SEQ ID NO:63, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pw460\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:64. Amino acids 17 to 29 of SEQ ID NO:64 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pw460\_5 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pw460\_5 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for pw460\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pw460\_5 demonstrated at least some similarity with sequences identified as AA447258 (zw93e03.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 784540 5', mRNA sequence), AA617801 (nq04f05.s1 NCI\_CGAP\_Lu1 Homo sapiens cDNA clone IMAGE 1142913), AC002486 (Human BAC clone RG367O17 from 7p15-p21, complete sequence), AC004837 (human genomic DNA fragments), and H45347 (yo65h03.r1 Homo sapiens cDNA clone 182837 5'). Based upon sequence similarity, pw460\_5 proteins and each similar protein or peptide may share at least some activity.

Clone "qa136\_1"

A polynucleotide of the present invention has been identified as clone "qa136\_1". qa136\_1 was isolated from a human adult cartilage (chondrosarcoma HTB-94 line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on

the basis of computer analysis of the amino acid sequence of the encoded protein. qa136\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qa136\_1 protein").

The nucleotide sequence of qa136\_1 as presently determined is reported in SEQ ID NO:65. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qa136\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:66. Amino acids 15 to 27 of SEQ ID NO:66 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qa136\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qa136\_1 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for qa136\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qa136\_1 demonstrated at least some similarity with sequences identified as AA758023 (ah67g02.s1 Soares testis NHT Homo sapiens cDNA clone 1320722 3', mRNA sequence), R69911 (yi47c02.r1 Homo sapiens cDNA clone 142370 5'), and T21835 (Human gene signature HUMGS03376; standard; cDNA to mRNA). Based upon sequence similarity, qa136\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five additional potential transmembrane domains within the qa136\_1 protein sequence, centered around amino acids 59, 136, 171, 201, and 268 of SEQ ID NO:66, respectively.

qa136\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 24 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "qy1261\_2"

A polynucleotide of the present invention has been identified as clone "qy1261\_2". qy1261\_2 was isolated from a human adult blood (promyelocytic Leukemia HL-60) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein.

qy1261\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qy1261\_2 protein").

The nucleotide sequence of qy1261\_2 as presently determined is reported in SEQ ID NO:67, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qy1261\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:68. Amino acids 100 to 112 of SEQ ID NO:68 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 113. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qy1261\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qy1261\_2 should be approximately 2500 bp.

The nucleotide sequence disclosed herein for qy1261\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qy1261\_2 demonstrated at least some similarity with sequences identified as AA076472 (zm91b06.r1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone 545267 5'), AA115700 (zl87g10.r1 Stratagene colon (#937204) Homo sapiens cDNA clone 511650 5', mRNA sequence), and AA190522 (zp85e07.r1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 627012 5'). The predicted amino acid sequence disclosed herein for qy1261\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qy1261\_2 protein demonstrated at least some similarity to the sequence identified as U49082 (transporter protein [Homo sapiens]). Based upon sequence similarity, qy1261\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts ten additional potential transmembrane domains within the qy1261\_2 protein sequence, centered around amino acids 80, 157, 203, 227, 286, 322, 365, 403, 426, and 462 of SEQ ID NO:68. The nucleotide sequence of qy1261\_2 indicates that it may contain one or more Alu repeat sequences.

#### Clone "rd432\_4"

A polynucleotide of the present invention has been identified as clone "rd432\_4". rd432\_4 was isolated from a human kidney (293 embryonal carcinoma cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S.

Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rd432\_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rd432\_4 protein").

5 The nucleotide sequence of rd432\_4 as presently determined is reported in SEQ ID NO:69, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rd432\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:70.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
10 rd432\_4 should be approximately 2200 bp.

The nucleotide sequence disclosed herein for rd432\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rd432\_4 demonstrated at least some similarity with sequences identified as AA662913 (nu92b03.s1 NCI\_CGAP\_Pr22 Homo sapiens cDNA clone  
15 IMAGE:1218125, mRNA sequence). Based upon sequence similarity, rd432\_4 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the rd432\_4 protein sequence, which includes amino acids 102-122 of SEQ ID NO:70. The nucleotide sequence of rd432\_4 indicates that it may contain one or more Alu repetitive elements.

20 rd432\_4 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 18 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "rb789\_14"

25 A polynucleotide of the present invention has been identified as clone "rb789\_14". rb789\_14 was isolated from a human kidney (293 embryonal carcinoma line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rb789\_14  
30 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rb789\_14 protein").

The nucleotide sequence of rb789\_14 as presently determined is reported in SEQ ID NO:71, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rb789\_14 protein

corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:72. Amino acids 9 to 21 of SEQ ID NO:72 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
5 should the predicted leader/signal sequence not be separated from the remainder of the rb789\_14 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rb789\_14 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for rb789\_14 was searched against the  
10 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rb789\_14 demonstrated at least some similarity with sequences identified as AL008582 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 223H9; HTGS phase 1), AL022393 (Homo sapiens DNA sequence from P1 p373c6 on chromosome 6p21.31-21.33. Contains zinc finger proteins, pseudogenes, ESTs  
15 and STS), N28823 (yx71f11.r1 Homo sapiens cDNA clone 267213 5'), and Q60944 (Human brain Expressed Sequence Tag EST01025; standard; DNA). Based upon sequence similarity, rb789\_14 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the rb789\_14 protein sequence, one centered around  
20 amino acid 30 and another around amino acid 75 of SEQ ID NO:72.

#### Clone "yd137\_1"

A polynucleotide of the present invention has been identified as clone "yd137\_1". yd137\_1 was isolated from a human adult brain cDNA library and was identified as  
25 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yd137\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yd137\_1 protein").

The nucleotide sequence of yd137\_1 as presently determined is reported in SEQ ID NO:73, and includes a poly(A) tail. What applicants presently believe to be the proper  
30 reading frame and the predicted amino acid sequence of the yd137\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:74. Amino acids 27 to 39 of SEQ ID NO:74 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 40. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain

should the predicted leader/signal sequence not be separated from the remainder of the yd137\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yd137\_1 should be approximately 789 bp.

5       The nucleotide sequence disclosed herein for yd137\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. yd137\_1 demonstrated at least some similarity with sequences identified as AI015619 (ov29g02.x1 Soares\_testis\_NHT Homo sapiens cDNA clone IMAGE:1638770 3' similar to WP:C34B2.10 CE16898; mRNA sequence). The predicted  
10 amino acid sequence disclosed herein for yd137\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted yd137\_1 protein demonstrated at least some similarity to the sequence identified as AF043693 (Caenorhabditis elegans cosmid C34B2). Based upon sequence  
15 similarity, yd137\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the yd137\_1 protein sequence, one centered around amino acid 30 and another around amino acid 55 of SEQ ID NO:74.

#### Clone "yd218\_1"

20       A polynucleotide of the present invention has been identified as clone "yd218\_1". yd218\_1 was isolated from a human adult brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yd218\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yd218\_1 protein").

25       The nucleotide sequence of yd218\_1 as presently determined is reported in SEQ ID NO:75, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the yd218\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76. Amino acids 2 to 14 of SEQ ID NO:76 are a predicted leader/signal sequence, with the predicted  
30 mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the yd218\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yd218\_1 should be approximately 900 bp.

The nucleotide sequence disclosed herein for yd218\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. yd218\_1 demonstrated at least some similarity with sequences identified as AA402818 (zu55f06.s1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 741923 3', mRNA sequence) and AI150344 (qf35b11.x1 Soares\_testis\_NHT Homo sapiens cDNA clone IMAGE:1751997 3', mRNA sequence). Based upon sequence similarity, yd218\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the yd218\_1 protein sequence, one centered around amino acid 66 and another around amino acid 100 of SEQ ID NO:76.

yd218\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 15 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "ye11\_1"

A polynucleotide of the present invention has been identified as clone "ye11\_1". ye11\_1 was isolated from a humna fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ye11\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ye11\_1 protein").

The nucleotide sequence of ye11\_1 as presently determined is reported in SEQ ID NO:77, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ye11\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ye11\_1 should be approximately 2700 bp.

The nucleotide sequence disclosed herein for ye11\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ye11\_1 demonstrated at least some similarity with sequences identified as AC005082 (\*\* SEQUENCING IN PROGRESS \*\* Homo sapiens clone RG271G13; HTGS phase 1, 7 unordered pieces). The predicted amino acid sequence disclosed herein for ye11\_1 was searched against the GenPept and GeneSeq amino acid

sequence databases using the BLASTX search protocol. The predicted ye11\_1 protein demonstrated at least some similarity to sequences identified as AF059569 (actin binding protein MAYVEN [Homo sapiens]) and R94386 (Human neural cell protein marker RR/B). MAYVEN is an actin-binding protein expressed in brain. Hidden markov model  
5 analysis reveals the presence of a BTB (BR-c/Ttk) domain in the predicted ye11\_1 protein. BTB domains are characteristic of certain bacterial membrane transport proteins. The MAYVEN protein is thought to contain a similar BTB motif, an indication that ye11\_1 and MAYVEN may share a similar function. Based upon sequence similarity, ye11\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII  
10 computer program predicts two potential transmembrane domains within the ye11\_1 protein sequence, one centered around amino acid 20 and another around amino acid 480 of SEQ ID NO:78.

ye11\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 57 kDa was detected in conditioned medium and  
15 membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "ye72\_1"

A polynucleotide of the present invention has been identified as clone "ye72\_1". ye72\_1 was isolated from a human fetal brain cDNA library and was identified as  
20 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ye72\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ye72\_1 protein").

The nucleotide sequence of ye72\_1 as presently determined is reported in SEQ ID NO:79, and includes a poly(A) tail. What applicants presently believe to be the proper  
25 reading frame and the predicted amino acid sequence of the ye72\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:80. Amino acids 24 to 36 of SEQ ID NO:80 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 37. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
30 the predicted leader/signal sequence not be separated from the remainder of the ye72\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ye72\_1 should be approximately 2261 bp.



The nucleotide sequence disclosed herein for ye72\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ye72\_1 demonstrated at least some similarity with sequences identified as AA968450 (op49d06.s1 Soares\_NFL\_T\_GBC\_S1 Homo sapiens cDNA clone  
5 IMAGE:1580171 3', mRNA sequence). The predicted amino acid sequence disclosed herein for ye72\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ye72\_1 protein demonstrated at least some similarity to sequences identified as U16258 (I kappa BR [Homo sapiens]) and W15483 (Human P28). Based upon sequence similarity, ye72\_1 proteins and each  
10 similar protein or peptide may share at least some activity. Hidden markov model analysis reveals the presence of three ankyrin repeats in the predicted ye72\_1 protein at amino acids 273 to 306, 307 to 339, and 341 to 373 of SEQ ID NO:80. The ankyrin 33-residue repeating motif, an L-shaped structure with protruding beta-hairpin tips, mediates specific macromolecular interactions with cytoskeletal, membrane, and regulatory proteins. The  
15 TopPredII computer program predicts an additional potential transmembrane domain within the ye72\_1 protein sequence centered around amino acid 140 of SEQ ID NO:80.

#### Clone "ye78\_1"

A polynucleotide of the present invention has been identified as clone "ye78\_1".  
20 ye78\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ye78\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ye78\_1 protein").

The nucleotide sequence of ye78\_1 as presently determined is reported in SEQ ID  
25 NO:81, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ye78\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:82. Amino acids 78 to 90 of SEQ ID NO:82 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 91. Amino acids 42 to 54 are also a possible  
30 leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 55. Due to the hydrophobic nature of leader/signal sequences, both of these predicted and possible leader sequences are likely to act as a transmembrane domain should either of them not be separated from the remainder of the ye78\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ye78\_1 should be approximately 2654 bp.

The nucleotide sequence disclosed herein for ye78\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ye78\_1 demonstrated at least some similarity with sequences identified as AA522797 (ni40c10.s1 NCI\_CGAP\_Lu1 Homo sapiens cDNA clone IMAGE:979314, mRNA sequence). Based upon sequence similarity, ye78\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four potential transmembrane domains within the ye78\_1 protein sequence, centered around amino acids 55, 75, 84, and 480 of SEQ ID NO:12, respectively.

#### Clone "ye90\_1"

A polynucleotide of the present invention has been identified as clone "ye90\_1". ye90\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ye90\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ye90\_1 protein").

The nucleotide sequence of ye90\_1 as presently determined is reported in SEQ ID NO:83, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ye90\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:84. Amino acids 7 to 19 of SEQ ID NO:84 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ye90\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ye90\_1 should be approximately 1505 bp.

The nucleotide sequence disclosed herein for ye90\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ye90\_1 demonstrated at least some similarity with sequences identified as AI079268 (oz32f06.x1 Soares\_total\_fetus\_Nb2HF8\_9w Homo sapiens cDNA clone IMAGE:1677059 3', mRNA sequence) and T25543 (Human gene signature HUMGS07715, standard; cDNA to mRNA). Based upon sequence similarity, ye90\_1

proteins and each similar protein or peptide may share at least some activity. Motifs analysis reveals the presence of a neutral zinc metallopeptidases, zinc-binding region signature beginning around amino acid residue 236 of SEQ ID NO:84; some known secreted proteins have this motif. The TopPredII computer program predicts two  
5 additional potential transmembrane domains within the ye90\_1 protein sequence, one centred around amino acid 195 and another around amino acid 300 of SEQ ID NO:84.

#### Clone "yi62\_1"

A polynucleotide of the present invention has been identified as clone "yi62\_1".  
10 yi62\_1 was isolated from a human adult brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yi62\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yi62\_1 protein").

The nucleotide sequence of yi62\_1 as presently determined is reported in SEQ ID  
15 NO:85, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the yi62\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:86. Amino acids 2 to 14 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal  
20 sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the yi62\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yi62\_1 should be approximately 1240 bp.

The nucleotide sequence disclosed herein for yi62\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. yi62\_1 demonstrated at least some similarity with sequences identified as R57572 (F3589 Fetal heart Homo sapiens cDNA clone F3589 5' end, mRNA sequence). Based upon sequence similarity, yi62\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four  
30 potential transmembrane domains within the yi62\_1 protein sequence, centered around amino acids 15, 75, 100, and 125 of SEQ ID NO:86, respectively.

Clone "yk78\_1"

A polynucleotide of the present invention has been identified as clone "yk78\_1". yk78\_1 was isolated from a human adult thymus cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yk78\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yk78\_1 protein").

The nucleotide sequence of yk78\_1 as presently determined is reported in SEQ ID NO:87, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the yk78\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:88. Amino acids 57 to 69 of SEQ ID NO:88 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 70. Amino acids 7 to 19 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of leader/signal sequences, both of these predicted and possible leader sequences are likely to act as a transmembrane domain should either of them not be separated from the remainder of the yk78\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yk78\_1 should be approximately 1088 bp.

The nucleotide sequence disclosed herein for yk78\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. yk78\_1 demonstrated at least some similarity with sequences identified as AC004921 (\*\* SEQUENCING IN PROGRESS \*\* Homo sapiens clone DJ0899E09; HTGS phase 1, 11 unordered pieces). Based upon sequence similarity, yk78\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the yk78\_1 protein sequence, one centered around amino acid 20 and another around amino acids 60 of SEQ ID NO:88.

Clone "yk251\_1"

A polynucleotide of the present invention has been identified as clone "yk251\_1". yk251\_1 was isolated from a human adult thymus cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yk251\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yk251\_1 protein").

The nucleotide sequence of yk251\_1 as presently determined is reported in SEQ ID NO:89, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the yk251\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:90. Amino acids 17 to 29 of SEQ ID NO:90 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the yk251\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yk251\_1 should be approximately 2558 bp.

The nucleotide sequence disclosed herein for yk251\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the databases. The TopPredII computer program predicts a potential transmembrane domain within the yk251\_1 protein sequence centered, around amino acid 20 of SEQ ID NO:90. The nucleotide sequence of yk251\_1 indicates that it may contain Alu and SVA repetitive elements.

#### Clone "yt14\_1"

A polynucleotide of the present invention has been identified as clone "yt14\_1". yt14\_1 was isolated from a human adult retina (WERI-Rb1 retinoblastoma line) cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. yt14\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "yt14\_1 protein").

The nucleotide sequence of yt14\_1 as presently determined is reported in SEQ ID NO:91, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the yt14\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:92. Amino acids 1 to 9 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 10. Due to the hydrophobic nature of this possible leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the yk251\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone yt14\_1 should be approximately 2429 bp.

The nucleotide sequence disclosed herein for yt14\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. yt14\_1 demonstrated at least some similarity with sequences identified as W07167 (za93b12.r1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 300095 5', mRNA sequence). The predicted amino acid sequence disclosed herein for yt14\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted yt14\_1 protein demonstrated at least some similarity to the sequence identified as AF002196 (weak similarity to Bacillus and Pseudomonas probable glucarate transporters (GI 709999 and PIR S27616) [Caenorhabditis elegans]). Based upon sequence similarity, yt14\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts six potential transmembrane domains within the yt14\_1 protein sequence, centered around amino acids 10, 40, 65, 90, 130, and 160 of SEQ ID NO:92, respectively. The nucleotide sequence of yt14\_1 indicates that it may contain Alu and L1 repetitive elements.

#### Clone "bf157\_16"

A polynucleotide of the present invention has been identified as clone "bf157\_16". bf157\_16 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bf157\_16 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "bf157\_16 protein").

The nucleotide sequence of bf157\_16 as presently determined is reported in SEQ ID NO:93, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bf157\_16 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:94.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bf157\_16 should be approximately 3480 bp.

The nucleotide sequence disclosed herein for bf157\_16 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bf157\_16 demonstrated at least some similarity with sequences identified as AA186595 (zo71g04.r1 Stratagene pancreas (#937208) Homo sapiens cDNA

clone 592374 5' similar to WP C16A3.3 CE04004 HUMAN ALPHA-FETOPROTEIN ENHANCER-BINDING PROTEIN), AA630405 (ac09b05.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 855921 3' similar to WP C16A3.3 CE04004 HUMAN ALPHA-FETOPROTEIN ENHANCER-BINDING PROTEIN; mRNA sequence), AF075104  
5 (Homo sapiens full length insert cDNA YR39H06), H49655 (yq20h07.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 274428 3'), Z28494 (H. sapiens partial cDNA sequence; clone 22G07; version 1; strand(-), single read), Z56794 (H.sapiens CpG island DNA genomic MseI fragment, clone), and Z64553 (H.sapiens CpG island DNA genomic MseI fragment, clone 139f5, forward read cpg139f5.ft1a). The predicted amino acid  
10 sequence disclosed herein for bf157\_16 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bf157\_16 protein demonstrated at least some similarity to sequences identified as R23962 (AFP-1. DNA encoding protein binding to alpha-fetoprotein gene enhancer -useful for prodn. of biological active protein), and U41534 (similar to yeast hypothetical protein  
15 (SP:YB9M\_YEAST,P38344); similar to human alpha-fetoprotein enhancer-binding protein (PIR:A41948) [Caenorhabditis elegans]). Based upon sequence similarity, bf157\_16 proteins and each similar protein or peptide may share at least some activity. Hidden Markov model and motifs analyses have revealed the presence of the following protein domains in the predicted bf157\_16 protein: four Zinc finger, C2H2 type, domains at  
20 amino acids 4 to 28, 67 to 91, 252 to 275, and 303 to 330 of SEQ ID NO:94; and a D-isomer-specific 2-hydroxyacid dehydrogenases signature at residues 119 to 131 of SEQ ID NO:94. A number of NAD-dependent 2-hydroxyacid dehydrogenases, with at least some specificity for the D-isomer of their substrate, have been shown to be functionally and structurally related. Clone bf157\_16 appears to encode a novel protein which may have  
25 NAD-dependent 2-hydroxyacid dehydrogenase activity.

bf157\_16 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 16 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

30

#### Clone "bk343\_2"

A polynucleotide of the present invention has been identified as clone "bk343\_2". bk343\_2 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer

analysis of the amino acid sequence of the encoded protein. bk343\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "bk343\_2 protein").

The nucleotide sequence of bk343\_2 as presently determined is reported in SEQ ID NO:95, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bk343\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:96.

Another possible reading frame within the bk343\_2 clone extends from nucleotide 45 to nucleotide 188 of SEQ ID NO:95, and encodes the amino acid sequence reported in SEQ ID NO:239. Amino acids 5 to 17 of SEQ ID NO:239 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:239.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bk343\_2 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for bk343\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bk343\_2 demonstrated at least some similarity with sequences identified as AA156969 (zo51d05.r1 Stratagene endothelial cell 937223 Homo sapiens cDNA clone 590409 5'), AA947938 (oe60c08.s1 NCI\_CGAP\_Lu5 Homo sapiens cDNA clone IMAGE:1416014 3', mRNA sequence), N31147 (yx52g05.r1 Homo sapiens cDNA clone 265400 5'), N42759 (yy22a09.r1 Homo sapiens cDNA clone 271960 5'), N47537 (yy90h10.s1 Homo sapiens cDNA clone 280867 3'), R68913 (yi43b04.r1 Homo sapiens cDNA clone 141967 5'), T24885 (Human gene signature HUMGS06991; standard; cDNA to mRNA), and T30099 (EST112339 Homo sapiens cDNA 5' end similar to None). The predicted amino acid sequence disclosed herein for bk343\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bk343\_2 protein demonstrated at least some similarity to sequences identified as Z72508 (F28H7.4 [Caenorhabditis elegans]) and Z78417 (C35C5.3 [Caenorhabditis elegans]). Based upon sequence similarity, bk343\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the bk343\_2 protein sequence centered around amino acid 36 of SEQ ID NO:96.



Clone "cd205\_2"

A polynucleotide of the present invention has been identified as clone "cd205\_2". cd205\_2 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cd205\_2 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "cd205\_2 protein").

The nucleotide sequence of cd205\_2 as presently determined is reported in SEQ ID NO:97, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cd205\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:98. Amino acids 92 to 104 of SEQ ID NO:98 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 105. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the cd205\_2 protein.

Another possible reading frame within the cd205\_2 clone extends from nucleotide 59 to nucleotide 478 of SEQ ID NO:97, and encodes the amino acid sequence reported in SEQ ID NO:240. The open reading frames encoding the amino acid sequences of SEQ ID NO:98 and SEQ ID NO:240 could be joined if one or more frame shifts were made in the nucleotide sequence of SEQ ID NO:97.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cd205\_2 should be approximately 1300 bp.

The nucleotide sequence disclosed herein for cd205\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cd205\_2 demonstrated at least some similarity with sequences identified as AA053543 (zl71f10.r1 Stratagene colon (#937204) Homo sapiens cDNA clone 510091 5' similar to gb:M77830 DESMOPLAKIN I AND II (HUMAN)), AC005332 (Homo sapiens chromosome 17, clone hRPK.147\_L\_13, complete sequence), N84944 (J1677F Homo sapiens cDNA clone J1677 5' similar to CHROMOSOME 4 (CLONE P4-661) STS4-563), N86274 (J7481F Fetal heart, Lambda ZAP Express Homo sapiens cDNA clone J7481 5' similar to CHROMOSOME 4 (CLONE P4-661) STS4-563), W68823 (zd37f04.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 342847 5', mRNA sequence), and Z54387 (H.sapiens CpG island DNA genomic MseI fragment, clone 10g3, reverse read cpg10g3.rt1a). Based upon sequence similarity, cd205\_2 proteins and each similar protein

or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the cd205\_2 protein sequence located around amino acid 105 of SEQ ID NO:98.

5        Clone "cw1292\_8"

A polynucleotide of the present invention has been identified as clone "cw1292\_8". cw1292\_8 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cw1292\_8 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cw1292\_8 protein").

The nucleotide sequence of cw1292\_8 as presently determined is reported in SEQ ID NO:99, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cw1292\_8 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:100. Amino acids 18 to 30 of SEQ ID NO:100 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 31. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the cw1292\_8 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw1292\_8 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for cw1292\_8 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw1292\_8 demonstrated at least some similarity with sequences identified as AA017976 (mh46h10.r1 Soares mouse placenta 4NbMP13.5 14.5 Mus), AA423855 (zv79c04.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 759846 3'), AA626784 (ad09f08.s1 Soares NbHFB Homo sapiens cDNA clone 877767 3', mRNA sequence), H23387 (ym57f05.r1 Homo sapiens cDNA clone 52337 5'), H78534 (yu13d06.r1 Homo sapiens cDNA clone 233675 5'), H79021 (yu13d06.s1 Homo sapiens cDNA clone 233675 3'), R44807 (yg23g06.s1 Homo sapiens cDNA clone 33217 3'), T24772 (Human gene signature HUMGS06848; standard; cDNA to mRNA), T97424 (ye53h08.r1 Homo sapiens cDNA clone 121503 5'), and Z44597 (H. sapiens partial cDNA sequence; clone c-25a05).

The predicted amino acid sequence disclosed herein for cw1292\_8 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cw1292\_8 protein demonstrated at least some similarity to the sequence identified as M33521 (HLA-B-associated transcript 3 (BAT3) [Homo]). Based upon sequence similarity, cw1292\_8 proteins and each similar protein or peptide may share at least some activity.

### Clone "cw1475\_2"

A polynucleotide of the present invention has been identified as clone "cw1475\_2". cw1475\_2 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cw1475\_2 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "cw1475\_2 protein").

The nucleotide sequence of cw1475\_2 as presently determined is reported in SEQ ID NO:101, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cw1475\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:102.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw1475\_2 should be approximately 2800 bp.

The nucleotide sequence disclosed herein for cw1475\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw1475\_2 demonstrated at least some similarity with sequences identified as AA527429 (ng41a10.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA clone IMAGE:937338, mRNA sequence), AD000092 (Homo sapiens DNA from chromosome 19p13.2 cosmids R31240, R30272 and R28549 containing the EKLF, GCDH, CRTCL, and RAD23A genes, genomic sequence), H98508 (yv90f08.r1 Homo sapiens cDNA clone 250023 5'), N25554 (yx76f08.s1 Homo sapiens cDNA clone 267687 3'), N50970 (yy94b06.s1 Homo sapiens cDNA clone 281171 3'), N81188 (yw36g06.r1 Homo sapiens cDNA clone 254362 5'), R32569 (yh54g03.r1 Homo sapiens cDNA clone 133588 5'), R81017 (yi94g02.r1 Homo sapiens cDNA clone 146930 5' similar to contains Alu repetitive element; contains MER30 repetitive element), T06537 (EST04426 Homo sapiens cDNA clone HFB DU83 similar to EST containing Alu repeat), T31594 (Probe (BLUR11) for Alu repeat sequence), and W30895 (zb78e12.r1 Soares senescent fibroblasts NbHSF Homo). Based upon sequence similarity, cw1475\_2 proteins and each similar protein or peptide may share at

least some activity. The nucleotide sequence of cw1475\_2 indicates that it may contain one or more of the following repetitive elements: Alu, SVA.

Clone "dd428\_4"

5 A polynucleotide of the present invention has been identified as clone "dd428\_4". dd428\_4 was isolated from a human adult testes cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dd428\_4 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "dd428\_4 protein").

10 The nucleotide sequence of dd428\_4 as presently determined is reported in SEQ ID NO:103, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dd428\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:104.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
15 dd428\_4 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for dd428\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dd428\_4 demonstrated at least some similarity with sequences identified as AC000057 (Human BAC clone RG067M09 from 7q21-7q22; HTGS phase 3, complete sequence), AC005500 (complete sequence), L27428 (Human L1 putative reverse transcriptase gene insertion in hamster, 3'end), T86176 (yd78c11.s1 Homo sapiens cDNA clone 114356 3' similar to gb L25879 EPOXIDE HYDROLASE (HUMAN); contains L1 repetitive element), X61307 (Staphylococcus aureus spa gene for protein A), and Z69647 (Human DNA sequence from cosmid E118G4, maps to 10cen and 11q13-q14). Based upon  
20 sequence similarity, dd428\_4 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of dd428\_4 indicates that it may contain L1 repeat sequences.

Clone "dh1073\_12"

30 A polynucleotide of the present invention has been identified as clone "dh1073\_12". dh1073\_12 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dh1073\_12 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "dh1073\_12 protein").

The nucleotide sequence of dh1073\_12 as presently determined is reported in SEQ ID NO:105, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dh1073\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:106.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dh1073\_12 should be approximately 2400 bp.

The nucleotide sequence disclosed herein for dh1073\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dh1073\_12 demonstrated at least some similarity with sequences  
10 identified as AA257983 (zs35h03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 687221 3' similar to TR G666014 G666014 SA SA GENE PRODUCT, COMPLETE CDS PRECURSOR; mRNA sequence), AA526325 (ni59g06.s1 NCI\_CGAP\_Ov2 Homo sapiens cDNA clone 981178 similar to contains Alu repetitive element), AF001549 (Human Chromosome 16 BAC clone CIT987SK-A-270G1, complete sequence), N57823 (yv59e04.s1  
15 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 247038 3'), and N68408 (za13c05.s1 Homo sapiens cDNA clone 292424 3'). The predicted amino acid sequence disclosed herein for cw1292\_8 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dh1073\_12 protein demonstrated at least some similarity to the sequence identified as AC003034 (Gene with  
20 similarity to rat kidney-specific (KS) gene [Homo sapiens]). Based upon sequence similarity, dh1073\_12 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of dh1073\_12 indicates that it may contain an Alu repetitive element.

25 Clone "dw78\_1"

A polynucleotide of the present invention has been identified as clone "dw78\_1". dw78\_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
30 analysis of the amino acid sequence of the encoded protein. dw78\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dw78\_1 protein").

The nucleotide sequence of dw78\_1 as presently determined is reported in SEQ ID NO:107, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the dw78\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:108.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dw78\_1 should be approximately 1400 bp.

5 The nucleotide sequence disclosed herein for dw78\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dw78\_1 demonstrated at least some similarity with sequences identified as AA807622 (nv65g11.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 1234724, mRNA sequence), AF086326 (Homo sapiens full length insert cDNA clone ZD54A02), D37980 (Dictyostelium discoideum DDCOF1 gene for cofilin, complete cds (exon1-2)), H26207 (yl53c04.r1 Homo sapiens cDNA clone 161958 5'), N72717 (za47h03.s1 Homo sapiens cDNA clone 295733 3' similar to contains Alu repetitive element; contains element L1 repetitive element), T23963 (Human gene signature HUMGS05917; standard; cDNA to mRNA), U14567 (\*\*ALU WARNING Human Alu-J subfamily consensus sequence), U43572 (Human alpha-N-acetylglucosaminidase (NAGLU) gene, complete cds), W42787 (zc25a04.s1 Soares senescent fibroblasts NbHSF Homo sapiens cDNA clone 323310 3'), and W73472 (zd54a02.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344426 3', mRNA sequence). The predicted amino acid sequence disclosed herein for dw78\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dw78\_1 protein demonstrated at least some similarity to the sequence identified as D32202 (alpha 1C adrenergic receptor isoform 2 [Homo sapiens]). Based upon sequence similarity, dw78\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the dw78\_1 protein sequence, one centered around amino acid 45 and another around amino acid 93 of SEQ ID NO:108. The nucleotide sequence of dw78\_1 indicates that it may contain an Alu repetitive element.

#### Clone "fh116\_11"

30 A polynucleotide of the present invention has been identified as clone "fh116\_11". fh116\_11 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. fh116\_11 is a full-length

clone, including the entire coding sequence of a secreted protein (also referred to herein as "fh116\_11 protein").

The nucleotide sequence of fh116\_11 as presently determined is reported in SEQ ID NO:109, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the fh116\_11 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:110.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone fh116\_11 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for fh116\_11 was searched against the  
10 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fh116\_11 demonstrated at least some similarity with sequences identified as AA054185 (zf51c06.r1 Soares retina N2b4HR Homo sapiens cDNA clone 380458 5'), AA057975 (mj57b02.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 480171 5' similar to WP:F57A8.2 CE05983), AA128902 (zn90a05.s1 Stratagene  
15 lung carcinoma 937218 Homo sapiens cDNA clone 565424 3'), AA426021 (zw49h09.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773441 3'), AA505926 (nh98g03.s1 NCI\_CGAP\_Br2 Homo sapiens cDNA clone 966580), AI079540 (oz04e08.x1 Soares\_fetal\_liver\_spleen\_1NFLS\_S1 Homo sapiens cDNA clone IMAGE:1674374 3' similar to WP:F57A8.2 CE05983; mRNA sequence), H68794 (yr91h09.s1 Homo sapiens  
20 cDNA clone 212705 3'), H86659 (yt02c04.r1 Homo sapiens cDNA clone 223110 5'), T24554 (Human gene signature HUMGS06604; standard; cDNA to mRNA), U96490 (Rattus norvegicus liver mRNA, complete cds), and W00635 (yy71d12.r1 Homo sapiens cDNA clone 278999 5' similar to contains element PTR5 repetitive element). The predicted amino acid sequence disclosed herein for fh116\_11 was searched against the GenPept and  
25 GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted fh116\_11 protein demonstrated at least some similarity to sequences identified as AF004876 (54TMp [Homo sapiens]), U96490 (unknown [Rattus norvegicus]), and Z70781 (F57A8.2 [Caenorhabditis elegans]). Based upon sequence similarity, fh116\_11 proteins and each similar protein or peptide may share at least some activity. The  
30 TopPredII computer program predicts five potential transmembrane domains within the fh116\_11 protein sequence, centered around amino acids 35 to 49, 136, 171, 215, and 270 of SEQ ID NO:110, respectively.

fh116\_11 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 28 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5        Clone "fy356\_14"

A polynucleotide of the present invention has been identified as clone "fy356\_14". fy356\_14 was isolated from a human fetal placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. fy356\_14 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "fy356\_14 protein").

The nucleotide sequence of fy356\_14 as presently determined is reported in SEQ ID NO:111, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the fy356\_14 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:112. Amino acids 385 to 397 of SEQ ID NO:112 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 398. Due to the hydrophobic nature of this possible leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the fy356\_14 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone fy356\_14 should be approximately 3700 bp.

The nucleotide sequence disclosed herein for fy356\_14 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fy356\_14 demonstrated at least some similarity with sequences identified as AA017639 (ze38c05.r1 Soares retina N2b4HR Homo sapiens cDNA clone 361256 5' similar to PIR S55385 S55385 PEA-15 protein - mouse), AA181529 (zp51f07.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 612997 3'), AA687129 (nv63d03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 1234469), AA811277 (ob68e06.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1336546, mRNA sequence), N53623 (yz04e01.r1 Homo sapiens cDNA clone 282072 5'), T25935 (Human gene signature HUMGS08167; standard; cDNA to mRNA), T24538 (Human gene signature HUMGS06585; standard; cDNA to mRNA), and X86809 (H.sapiens mRNA for



major astrocytic phosphoprotein PEA-15). The predicted amino acid sequence disclosed herein for fy356\_14 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted fy356\_14 protein demonstrated at least some similarity to the sequence identified as X86809 (PEA-15 gene product [Homo sapiens]). Based upon sequence similarity, fy356\_14 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the fy356\_14 protein sequence, centered around amino acid 398 of SEQ ID NO:112.

10      Clone "iw66\_1"

A polynucleotide of the present invention has been identified as clone "iw66\_1". iw66\_1 was isolated from a human adult retina (WERI-Rb1 retinoblastoma line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. iw66\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "iw66\_1 protein").

The nucleotide sequence of iw66\_1 as presently determined is reported in SEQ ID NO:113, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the iw66\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:114. Amino acids 9 to 21 of SEQ ID NO:114 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the iw66\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone iw66\_1 should be approximately 1450 bp.

The nucleotide sequence disclosed herein for iw66\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. iw66\_1 demonstrated at least some similarity with sequences identified as AA216917 (mv75h11.r1 Soares mouse 3NME12 5 Mus musculus cDNA clone 660933 5'), AA339406 (EST44484 Fetal brain I Homo sapiens cDNA 5' end), AI275861 (ql68b12.x1 Soares\_NhHMPu\_S1 Homo sapiens cDNA clone IMAGE:1877471 3', mRNA

sequence), Q61257 (Human brain Expressed Sequence Tag EST01278; standard; DNA), R89651 (ym97c08.r1 Homo sapiens cDNA clone 166862 5'), W53584 (md55f06.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 372323 5'), and Z60886 (H.sapiens CpG island DNA genomic MseI fragment, clone 38a8, reverse read cpg38a8.rt1a). The predicted amino acid sequence disclosed herein for iw66\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted iw66\_1 protein demonstrated at least some similarity to sequences identified as AF004874 (latent TGF-beta binding protein-2 [Mus musculus]), L29029 (amino acid feature Rod protein domain, aa 266 468; amino acid feature globular protein domain, aa 32 .. 265 [Chlamydomonas reinhardtii]), R27150 (PspA fragment), and R79478 (Mouse LTBP-2). Based upon sequence similarity, iw66\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the iw66\_1 protein sequence, centered around amino acids 45, 74, and 158 of SEQ ID NO:114, respectively. The nucleotide sequence of iw66\_1 indicates that it may contain one or more of the following repetitive elements: MIR.

#### Clone "kh13\_4"

A polynucleotide of the present invention has been identified as clone "kh13\_4". kh13\_4 was isolated from a human adult testes cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. kh13\_4 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "kh13\_4 protein").

The nucleotide sequence of kh13\_4 as presently determined is reported in SEQ ID NO:115, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the kh13\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:116.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone kh13\_4 should be approximately 950 bp.

The nucleotide sequence disclosed herein for kh13\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. kh13\_4 demonstrated at least some similarity with sequences identified as AA435981 (zu01f08.s1 Soares testis NHT Homo sapiens cDNA clone 730599 3'), AA436078 (zu01f08.r1 Soares testis NHT Homo sapiens cDNA clone 730599 5'),

AA778636 (af87c04.s1 Soares testis NHT Homo sapiens cDNA clone 1048998 3' similar to gb:M94856 PSORIASIS-ASSOCIATED FATTY ACID BINDING PROTEIN HOMOLOG (HUMAN); mRNA sequence), M94856 (Human fatty acid binding protein homologue (PA-FABP) mRNA, complete cds), and Q66842 (Melanogenic inhibitor; standard; DNA).

5 The predicted amino acid sequence disclosed herein for kh13\_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted kh13\_4 protein demonstrated at least some similarity to sequences identified as M94856 (fatty acid binding protein homologue [Homo sapiens]) and R55866 (Melanogenic inhibitor). Fatty acid binding protein homologue (M94856) is described as

10 "a novel keratinocyte protein (psoriasis-associated fatty acid-binding protein [PA-FABP]) that is highly up-regulated in psoriatic skin and that shares similarity to fatty acid-binding proteins." Based upon sequence similarity, kh13\_4 proteins and each similar protein or peptide may share at least some activity.

15 Clone "ko258\_4"

A polynucleotide of the present invention has been identified as clone "ko258\_4". ko258\_4 was isolated from a human adult uterus cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ko258\_4 is a full-length clone, including the entire coding sequence

20 of a novel protein (also referred to herein as "ko258\_4 protein").

The nucleotide sequence of ko258\_4 as presently determined is reported in SEQ ID NO:117, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ko258\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:118.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ko258\_4 should be approximately 2500 bp.

The nucleotide sequence disclosed herein for ko258\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ko258\_4 demonstrated at least some similarity with sequences

30 identified as AC002401 (\*\*SEQUENCING IN PROGRESS\*\*) Homo sapiens chromosome 17, clone RPC875H18; HTGS phase 1, 4 unordered pieces), AC002401 (Homo sapiens chromosome 17, clone RPC875H18, complete sequence), C15329 (Human fetal brain cDNA 5'-end GEN-133H10, mRNA sequence), AF035306 (Homo sapiens clone 23771 mRNA sequence), and R28382 (IMAGE 3p clone). Based upon sequence similarity,

ko258\_4 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the ko258\_4 protein sequence, centered around amino acid 28 of SEQ ID NO:118.

5        Clone "kv10\_8"

A polynucleotide of the present invention has been identified as clone "kv10\_8". kv10\_8 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. kv10\_8 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "kv10\_8 protein").

The nucleotide sequence of kv10\_8 as presently determined is reported in SEQ ID NO:119, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the kv10\_8 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:120.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone kv10\_8 should be approximately 4300 bp.

The nucleotide sequence disclosed herein for kv10\_8 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. kv10\_8 demonstrated at least some similarity with sequences identified as AA418842 (zw01e12.s1 Soares NhHMPu S1 Homo sapiens cDNA clone 768046 3'), AC004228 (\*\* SEQUENCING IN PROGRESS \*\* Homo sapiens Chromosome 11q12 pac pDJ519o3; HTGS phase 1, 18 unordered pieces), AF052108 (Homo sapiens clone 23687 mRNA sequence), R00761 (ye78b11.s1 Homo sapiens cDNA clone 123837 3'), T83434 (yd46b04.r1 Homo sapiens cDNA clone 111247 5'), T84080 (yd46b04.s1 Homo sapiens cDNA clone 111247 3'), and U00594 (Mustela vison unknown mRNA down regulated by TGF-beta, partial sequence). Based upon sequence similarity, kv10\_8 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the kv10\_8 protein sequence, centered around amino acids 35 to 45 of SEQ ID NO:120. The nucleotide sequence of kv10\_8 indicates that it may contain one or more of the following repetitive elements: Alu, SVA.

Clone "LL89\_3"

A polynucleotide of the present invention has been identified as clone "LL89\_3". LL89\_3 was isolated from a human adult thyroid cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. LL89\_3 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "LL89\_3 protein").

The nucleotide sequence of LL89\_3 as presently determined is reported in SEQ ID NO:121, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the LL89\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:122.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone LL89\_3 should be approximately 900 bp.

The nucleotide sequence disclosed herein for LL89\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. LL89\_3 demonstrated at least some similarity with sequences identified as AL031010 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 422F24, complete sequence), H78002 (yu82h09.r1 Homo sapiens cDNA clone 240353 5'), and W90018 (zh72c08.s1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 417614 3'). Based upon sequence similarity, LL89\_3 proteins and each similar protein or peptide may share at least some activity.

Clone "mc300\_1"

A polynucleotide of the present invention has been identified as clone "mc300\_1". mc300\_1 was isolated from a human adult thyroid cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. mc300\_1 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "mc300\_1 protein").

The nucleotide sequence of mc300\_1 as presently determined is reported in SEQ ID NO:123, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the mc300\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:124.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone mc300\_1 should be approximately 2600 bp.

The nucleotide sequence disclosed herein for mc300\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. mc300\_1 demonstrated at least some similarity with sequences identified as AA142942 (IMAGE 3p clone), AA315222 (EST187017 Colon carcinoma (HCC) cell line Homo sapiens cDNA 5' end), AA142942 (zl43c04.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 504678 3'), AI246503 (qn64a06.x1 NCI\_CGAP\_HN4 Homo sapiens cDNA clone IMAGE:1902994 3', mRNA sequence), D61461 (Human fetal brain cDNA 5'-end GEN-404B08), D79662 (Human aorta cDNA 5'-end GEN-300D05, mRNA sequence), H93575 (yv14h11.s1 Homo sapiens cDNA clone 242757 3'), T25928 (Human gene signature HUMGS08160; standard; cDNA to mRNA), and W93059 (zd93h06.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 357083 3'). Based upon sequence similarity, mc300\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of mc300\_1 indicates that it may contain one or more Alu repetitive elements.

15

#### Clone "ml227\_1"

A polynucleotide of the present invention has been identified as clone "ml227\_1". ml227\_1 was isolated from a human adult brain (caudate nucleus) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ml227\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ml227\_1 protein").

The nucleotide sequence of ml227\_1 as presently determined is reported in SEQ ID NO:125, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ml227\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:126.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ml227\_1 should be approximately 2700 bp.

30

The nucleotide sequence disclosed herein for ml227\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ml227\_1 demonstrated at least some similarity with sequences identified as AA857826 (oe88e05.s1 NCI\_CGAP\_Co12 Homo sapiens cDNA clone IMAGE:1418720 3', mRNA sequence), F18464 (H.sapiens EST sequence (017-T4-16) from

skeletal muscle), H30845 (yo78d11.r1 Homo sapiens cDNA clone 184053 5'), T06839 (EST04728 Homo sapiens cDNA clone HFBDZ66), T19759 (Human gene signature HUMGS00834), T26021 (Human gene signature HUMGS08257; standard; cDNA to mRNA), and Z69043 (H.sapiens mRNA translocon-associated protein delta subunit precursor). The predicted amino acid sequence disclosed herein for ml227\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ml227\_1 protein demonstrated at least some similarity to the sequence identified as Z69664 (K04D7.5 [Caenorhabditis elegans]). Based upon sequence similarity, ml227\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts six potential transmembrane domains within the ml227\_1 protein sequence, centered around amino acids 465, 510, 560, 572, 595, and 615 of SEQ ID NO:126, respectively.

#### Clone "mm367\_6"

A polynucleotide of the present invention has been identified as clone "mm367\_6". mm367\_6 was isolated from a human adult retina (WERI-Rb1 retinoblastoma line) cDNA library and was identified as encoding a protein. mm367\_6 is a full-length clone, including the entire coding sequence of a protein (also referred to herein as "mm367\_6 protein").

The nucleotide sequence of mm367\_6 as presently determined is reported in SEQ ID NO:127, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the mm367\_6 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:128.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone mm367\_6 should be approximately 2600 bp.

The nucleotide sequence disclosed herein for mm367\_6 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. mm367\_6 demonstrated at least some similarity with sequences identified as AA114127 (zn65f02.r1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 563067 5'), AA127284 (zn91c12.r1 Stratagene lung carcinoma 937218 Homo sapiens cDNA clone 565558 5'), AA173842 (zp30d01.r1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 610945 5'), AF000364 (Homo sapiens heterogeneous nuclear ribonucleoprotein R mRNA, complete CDs), N31934 (yy22d10.s1 Homo sapiens cDNA clone 271987 3'), T24354 (Human gene signature HUMGS06385; standard; cDNA to

mRNA), U48271 (*Dictyostelium discoideum* UbpA deubiquitinase mRNA, complete CDs), W16579 (zb13g11.r1 Soares fetal lung NbHL19W *Homo sapiens* cDNA clone 301988 5'), and W72461 (zd67f06.s1 Soares fetal heart NbHH19W *Homo sapiens* cDNA clone 345731 3'). The predicted amino acid sequence disclosed herein for mm367\_6 was searched  
5 against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted mm367\_6 protein demonstrated at least some similarity to sequences identified as AF000364 (heterogeneous nuclear ribonucleoprotein R [*Homo sapiens*]) and W26553 (Human heterogeneous nuclear ribonucleoprotein (hnRNP) A2). Based upon sequence similarity, mm367\_6 proteins and each similar protein or peptide  
10 may share at least some activity.

mm367\_6 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 79 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

15 Clone "mt124\_3"

A polynucleotide of the present invention has been identified as clone "mt124\_3". mt124\_3 was isolated from a human adult testes cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. mt124\_3 is a full-length clone, including the entire coding sequence  
20 of a novel protein (also referred to herein as "mt124\_3 protein").

The nucleotide sequence of mt124\_3 as presently determined is reported in SEQ ID NO:129, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the mt124\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:130.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone mt124\_3 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for mt124\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. mt124\_3 demonstrated at least some similarity with sequences  
30 identified as AA435386 (ve15h01.r1 Soares mouse NbMH *Mus musculus* cDNA clone 818257 5' similar to TR:E198756 E198756 PUTATIVE ORF), AI185116 (qe51g07.x1 Soares\_fetal\_lung\_NbHL19W *Homo sapiens* cDNA clone IMAGE 1742556 3' similar to TR Q92564 Q92564 MYELOBLAST KIAA0276 ; mRNA sequence), C03847 (Human Heart cDNA, clone 3NHC2256), N74186 (za76h03.s1 *Homo sapiens* cDNA clone 298517 3'),



T24234 (Human gene signature HUMGS06248; standard; cDNA to mRNA), W87997 (mf65b06.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 419123 5'), and Z86062 (Human DNA sequence from PAC 121G13 on chromosome 6 contains flow sorted chromosome 6 HindIII fragment ESTs, polymorphic CA repeat, CpG island, CpG island genomic fragments). The predicted amino acid sequence disclosed herein for mt124\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted mt124\_3 protein demonstrated at least some similarity to sequences identified as AL024499 (H38K22.2 [Caenorhabditis elegans]) and D87466 (Similar to S.cerevisiae hypothetical protein L3111 (S59316) [Homo sapiens]).

Based upon sequence similarity, mt124\_3 proteins and each similar protein or peptide may share at least some activity.

#### Clone "nf56\_3"

A polynucleotide of the present invention has been identified as clone "nf56\_3". nf56\_3 was isolated from a human adult brain (substantia nigra) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. nf56\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "nf56\_3 protein").

The nucleotide sequence of nf56\_3 as presently determined is reported in SEQ ID NO:131, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the nf56\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:132. Amino acids 3 to 15 of SEQ ID NO:132 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the nf56\_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone nf56\_3 should be approximately 5000 bp.

The nucleotide sequence disclosed herein for nf56\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. nf56\_3 demonstrated at least some similarity with sequences

identified as H08054 (yl86a09.s1 Homo sapiens cDNA clone 44915 3'), Q60495 (Human brain Expressed Sequence Tag EST02500; standard; cDNA), T25509 (Human gene signature HUMGS07678), W34534 (mc58h01.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 352753 5'), and Z64987 (H.sapiens CpG island DNA genomic MseI  
5 fragment, clone 186b1, reverse read cpg186b1.rt1b). The predicted amino acid sequence disclosed herein for nf56\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted nf56\_3 protein demonstrated at least some similarity to sequences identified as D86983 (similar to D.melanogaster peroxidase (U11052) [Homo sapiens]), R25079 (Drosophila SLIT protein  
10 involved in axon pathway development), and X53959 (slit protein [Drosophila melanogaster]). Based upon sequence similarity, nf56\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the nf56\_3 protein sequence, one centered around amino acid 514 and another around amino acid 628 of SEQ ID NO:132.

15

#### Clone "qy442\_2"

A polynucleotide of the present invention has been identified as clone "qy442\_2". qy442\_2 was isolated from a human adult blood (promyelocytic leukemia HL-60 line) cDNA library using methods which are selective for cDNAs encoding secreted proteins  
20 (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qy442\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qy442\_2 protein").

The nucleotide sequence of qy442\_2 as presently determined is reported in SEQ  
25 ID NO:133, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qy442\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:134. Amino acids 3 to 15 of SEQ ID NO:134 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the  
30 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qy442\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qy442\_2 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for qy442\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qy442\_2 demonstrated at least some similarity with sequences identified as AI081522 (on04e12.x1 NCI\_CGAP\_Kid3 Homo sapiens cDNA clone  
5 IMAGE:1555726 3' similar to contains Alu repetitive element; mRNA sequence) and AA449854 (zx37a06.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 788626 5'). Based upon sequence similarity, qy442\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the qy442\_2 protein sequence, centered around amino acid  
10 68 of SEQ ID NO:20. The nucleotide sequence of qy442\_2 indicates that it may contain one or more Alu repetitive elements.

#### Clone "rj214\_14"

A polynucleotide of the present invention has been identified as clone "rj214\_14".  
15 rj214\_14 was isolated from a human adult neural (neuroepithelioma HTB-10 line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rj214\_14 is a full-length clone, including the entire coding sequence of a secreted protein  
20 (also referred to herein as "rj214\_14 protein").

The nucleotide sequence of rj214\_14 as presently determined is reported in SEQ ID NO:135, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rj214\_14 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:136.  
25 Amino acids 3 to 15 of SEQ ID NO:136 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the rj214\_14 protein.

30 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rj214\_14 should be approximately 900 bp.

The nucleotide sequence disclosed herein for rj214\_14 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rj214\_14 demonstrated at least some similarity with sequences

identified as AA167035 (zp05c10.s1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone 595506 3' similar to TR:G563357 G563357 GENES RAS1, RLB1 AND RLC1; mRNA sequence), AA491109 (aa52d09.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 824561 5' similar to TR G563357 G563357 GENES RAS1, RLB1 AND RLC1), and  
5 AI189156 (qd04c02.x1 Soares\_placenta\_8to9weeks\_2NbHP8to9W Homo sapiens cDNA clone IMAGE:1722722 3' similar to TR:O01437 O01437 SIMILAR TO DROSOPHILA RLC1 GENE PRODUCT; mRNA sequence). The predicted amino acid sequence disclosed herein for rj214\_14 was searched against the GenPept and GeneSeq amino acid sequence  
10 databases using the BLASTX search protocol. The predicted rj214\_14 protein demonstrated at least some similarity to sequences identified as U97016 (similar to drosophila Rlc1 gene product (NID g563361) and S. cerevisiae mitochondrial 60S ribosomal protein L4 (YML4) (NID g459259) [Caenorhabditis elegans]), and X73219 (Rlc1). Drosophila Rlc1 is a basic protein that is bound to the inner face of the cell membrane. Transcription mapping and nucleotide sequence analysis reveal that Rlc1 lies in the same  
15 genomic region as Drosophila Ras1 and shows expression patterns that are similar to those of Ras1. It has been demonstrated (Ezer *et al.*, 1994, *Dev. Dyn.* 201(2): 179-190, which is incorporated by reference herein) that during embryogenesis Ras1 transcripts are restricted mainly to the embryonic central nervous system, suggesting that the Rlc1 gene product also may have a role in these nerve cells. Based upon sequence similarity,  
20 rj214\_14 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the rj214\_14 protein sequence, centered around amino acid 32 of SEQ ID NO:136.

rj214\_14 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 22 kDa was detected in membrane fractions using SDS  
25 polyacrylamide gel electrophoresis.

#### Clone "rk80\_3"

A polynucleotide of the present invention has been identified as clone "rk80\_3". rk80\_3 was isolated from a human adult tumor (colorectal adenocarcinoma SW480 line)  
30 cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. rk80\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "rk80\_3 protein").

The nucleotide sequence of rk80\_3 as presently determined is reported in SEQ ID NO:137, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the rk80\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:138. Amino acids 6 to 18 of SEQ ID NO:138 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the rk80\_3 protein.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone rk80\_3 should be approximately 1096 bp.

The nucleotide sequence disclosed herein for rk80\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. rk80\_3 demonstrated at least some similarity with sequences identified as AA418955 (zw01c10.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 15 768018 5', mRNA sequence), AB004061 (domestic pig mRNA for STAT2, complete CDs, a signal transducer and activator of transcription), C06368 (similar to none), and U38443 (Human clone JkA3 mRNA induced upon T-cell activation, 3' end). The predicted rk80\_3 protein demonstrated at least some similarity to granulocyte-colony stimulating factor (G-CSF) and interleukin-6 (IL-6). Hidden Markov model analysis has revealed the presence of an IL-6/G-CSF/mast cell growth factor (MGF) family signature at amino acids 69 to 181 of SEQ ID NO:138. This family of cytokines are glycoproteins of about 170 to 180 amino acid residues in size that contain four conserved cysteine residues involved in two disulfide bonds. rk80\_3 appears to encode a novel cytokine in the IL-6/G-CSF family. 20 Based upon sequence similarity, rk80\_3 proteins and each similar protein or peptide may share at least some activity.

rk80\_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 24 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

30

#### Clone "au36\_42"

A polynucleotide of the present invention has been identified as clone "au36\_42". au36\_42 was isolated from a human adult testes cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of

the encoded protein. au36\_42 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "au36\_42 protein").

The nucleotide sequence of au36\_42 as presently determined is reported in SEQ ID NO:139, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the au36\_42 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:140.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone au36\_42 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for au36\_42 was searched against the  
10 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No significant hits were found in the database. The nucleotide sequence of au36\_42 indicates that it may contain a L1ME repetitive element.

Clone "bo549\_13"

15 A polynucleotide of the present invention has been identified as clone "bo549\_13". bo549\_13 was isolated from a human adult retina cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bo549\_13 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "bo549\_13 protein").

20 The nucleotide sequence of bo549\_13 as presently determined is reported in SEQ ID NO:141, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bo549\_13 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:142. The region of SEQ ID NO:141 at nucleotides 518 and 519 may represent the border of an  
25 alternatively spliced exon.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bo549\_13 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for bo549\_13 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
30 FASTA search protocols. bo549\_13 demonstrated at least some similarity with sequences identified as AI261562 (qz30c06.x1 NCI\_CGAP\_Kid11 Homo sapiens cDNA clone IMAGE 2028394 3' similar to TR Q63061 Q63061 HYPOTHETICAL 4.7 KD PROTEIN; mRNA sequence) and J02649 (Rat stomach (H+,K+)-ATPase mRNA, complete cds). The predicted amino acid sequence disclosed herein for bo549\_13 was searched against the GenPept and

GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bo549\_13 protein demonstrated at least some similarity to sequences identified as J02649 (unknown protein [Rattus norvegicus]). Based upon sequence similarity, bo549\_13 proteins and each similar protein or peptide may share at least some activity.

5

Clone "da529\_3"

A polynucleotide of the present invention has been identified as clone "da529\_3". da529\_3 was isolated from a human fetal placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
10 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. da529\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "da529\_3 protein").

The nucleotide sequence of da529\_3 as presently determined is reported in SEQ  
15 ID NO:143, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the da529\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:144. Amino acids 59 to 71 of SEQ ID NO:144 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 72. Due to the  
20 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the da529\_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone da529\_3 should be approximately 1150 bp.

25 The nucleotide sequence disclosed herein for da529\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. da529\_3 demonstrated at least some similarity with sequences identified as AI189911 (qd33e06.x1 Soares\_placenta\_8to9weeks\_2NbHP8to9W Homo sapiens cDNA clone IMAGE 1725538 3' similar to TR O42204 O42204 PUTATIVE  
30 TRANSMEMBRANE PROTEIN E3-16; mRNA sequence), T35254 (EST82005 Homo sapiens cDNA 5' end similar to None), U76253 (Mus musculus E25B protein mRNA, complete cds), V43619 (Human secreted protein 19 encoding DNA), W28608 (49b1 Human retina cDNA randomly primed sublibrary Homo sapiens cDNA), and W41628 (mc47c10.r1 Soares mouse p3NMF19). The predicted amino acid sequence disclosed herein for

da529\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted da529\_3 protein demonstrated at least some similarity to sequences identified as AF03895 (E25 protein [Homo sapiens]) and W63699 (Human secreted protein 19). Based upon sequence similarity, da529\_3 proteins and each similar protein or peptide may share at least some activity.

#### Clone "dm365\_3"

A polynucleotide of the present invention has been identified as clone "dm365\_3". A cDNA clone was first isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate dm365\_3 from a human fetal brain cDNA library. dm365\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dm365\_3 protein").

The nucleotide sequence of dm365\_3 as presently determined is reported in SEQ ID NO:145, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dm365\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:146. Amino acids 1 to 13 of SEQ ID NO:146 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 14. Amino acids 40 to 52 of SEQ ID NO:146 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning in that case at amino acid 53. Due to the hydrophobic nature of each of these predicted leader/signal sequences, each predicted leader/signal sequence is likely to act as a transmembrane domain should it not be separated from the remainder of the dm365\_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dm365\_3 should be approximately 1300 bp.

The nucleotide sequence disclosed herein for dm365\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dm365\_3 demonstrated at least some similarity with sequences identified as AC005533 (\*\* SEQUENCING IN PROGRESS \*\* Homo sapiens clone DJ0794K21; HTGS phase 1, 22 unordered pieces), AI125562 (qd94d09.x1 Soares testis NHT Homo sapiens cDNA clone IMAGE 1737137 3', mRNA sequence), R02268 (ye85c10.r1



Homo sapiens cDNA clone 124530 5' similar to contains LTR5 repetitive element), and V90427 (EST clone DM365). Based upon sequence similarity, dm365\_3 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of dm365\_3 indicates that it may contain repetitive sequences.

- 5 dm365\_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 23 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "fa171\_1"

- 10 A polynucleotide of the present invention has been identified as clone "fa171\_1". fa171\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. fa171\_1 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "fa171\_1 protein").

- 15 The nucleotide sequence of fa171\_1 as presently determined is reported in SEQ ID NO:147, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the fa171\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:148.

- The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 fa171\_1 should be approximately 2500 bp.

- The nucleotide sequence disclosed herein for fa171\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fa171\_1 demonstrated at least some similarity with sequences identified as AA446057 (zw66d04.r1 Soares testis NHT Homo sapiens cDNA clone 781159  
25 5', mRNA sequence), AC002099 (\*\* SEQUENCING IN PROGRESS \*\* Genomic sequence from Human 9q34; HTGS phase 1, 2 unordered pieces), AC002355 (\*\* SEQUENCING IN PROGRESS \*\* Genomic sequence from Human 9q34; HTGS phase 1, 7 unordered pieces), and U10185 (Xenopus laevis XPMC2 protein mRNA, complete cds). The predicted amino acid sequence disclosed herein for fa171\_1 was searched against the GenPept and  
30 GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted fa171\_1 protein demonstrated at least some similarity to sequences identified as R67549 (Fruiting body inducing polypeptide) and U10185 (XPMC2 protein [Xenopus laevis]). XPMC2 is a Xenopus cDNA clone that can rescue several different yeast mitotic catastrophe mutants defective in Wee1 kinase function, and is a nuclear protein. Based

upon sequence similarity, fa171\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "lp572\_2"

5 A polynucleotide of the present invention has been identified as clone "lp572\_2". lp572\_2 was isolated from a human adult blood (peripheral blood mononuclear cells treated with granulocyte-colony stimulating factor *in vivo*) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or  
10 was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. lp572\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "lp572\_2 protein").

The nucleotide sequence of lp572\_2 as presently determined is reported in SEQ ID NO:149, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the lp572\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:150. Amino acids 79 to 91 of SEQ ID NO:150 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 92. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a  
20 transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the lp572\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone lp572\_2 should be approximately 2100 bp.

The nucleotide sequence disclosed herein for lp572\_2 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. lp572\_2 demonstrated at least some similarity with sequences identified as AA489012 (aa56a03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone 824908 3'), AA533633 (nf73b09.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA clone IMAGE 925529, mRNA sequence), AC004686 (Homo sapiens chromosome 17, clone hRPC.1073\_F\_15, complete sequence), T18977 (g07030t Testis 1 Homo sapiens cDNA clone g07030 5' end),  
30 T21490 (Human gene signature HUMGS02862), and W73324 (zd01h01.r1 Pancreatic Islet Homo sapiens cDNA clone 339409 5'). The predicted amino acid sequence disclosed herein for lp572\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted lp572\_2 protein

demonstrated at least some similarity to sequences identified as AL03262 (predicted using Genefinder [Caenorhabditis elegans]). Based upon sequence similarity, lp572\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five additional potential transmembrane domains within the  
5 lp572\_2 protein sequence, centered around amino acids 129, 263, 286, 326, and 378 of SEQ ID NO:150, respectively.

#### Clone "pe246\_1"

A polynucleotide of the present invention has been identified as clone "pe246\_1".  
10 pe246\_1 was isolated from a human adult blood (chronic myelogenous leukemia line K562) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pe246\_1 is a full-length clone, including the entire coding sequence  
15 of a secreted protein (also referred to herein as "pe246\_1 protein").

The nucleotide sequence of pe246\_1 as presently determined is reported in SEQ ID NO:151, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pe246\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:152.  
20 Amino acids 193 to 205 of SEQ ID NO:152 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 206. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pe246\_1 protein.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pe246\_1 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for pe246\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pe246\_1 demonstrated at least some similarity with sequences  
30 identified as AA234138 (zr51b06.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 666899 5' similar to SW FCEB\_HUMAN Q01362 HIGH AFFINITY IMMUNOGLOBULIN EPSILON RECEPTOR BETA-SUBUNIT), AA418443 (zv92e05.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 767264 5' similar to SW FCEB\_RAT P13386 HIGH AFFINITY IMMUNOGLOBULIN EPSILON RECEPTOR BETA-SUBUNIT; mRNA sequence),

AC004584 (Homo sapiens chromosome 17, clone hRPC1107\_A\_17, complete sequence), M74509 (Human endogenous retrovirus type C oncovirus sequence), and V57903 (Hereditary haemochromatosis subregion from an HH affected individual). The predicted amino acid sequence disclosed herein for pe246\_1 was searched against the GenPept and  
5 GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pe246\_1 protein demonstrated at least some similarity to sequences identified as L35848 (IgE receptor beta subunit [Homo sapiens]), R05026 (Beta subunit of rat high affinity IgE receptor Fc(epsilon)RI), and R42341 (Subunit of the human IgE receptor). The first 359 nucleotides of SEQ ID NO:13 is similar in sequence to that of M74509 (Human  
10 endogenous retrovirus type C oncovirus sequence) and also to several genomic sequences as a result. It appears that this region may be retroviral DNA that has been incorporated into the genome. Based upon sequence similarity, pe246\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the pe246\_1 protein  
15 sequence, centered around amino acids 86, 115, and 154 of SEQ ID NO:152, respectively.

#### Clone "qf122\_3"

A polynucleotide of the present invention has been identified as clone "qf122\_3". qf122\_3 was isolated from a human adult bladder (carcinoma line 5637) cDNA library and  
20 was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qf122\_3 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "qf122\_3 protein").

The nucleotide sequence of qf122\_3 as presently determined is reported in SEQ ID NO:153. What applicants presently believe to be the proper reading frame and the  
25 predicted amino acid sequence of the qf122\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:154.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qf122\_3 should be approximately 1700 bp.

The nucleotide sequence disclosed herein for qf122\_3 was searched against the  
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qf122\_3 demonstrated at least some similarity with sequences identified as AA206909 (zq80d10.r1 Stratagene hNT neuron (#937233) Homo sapiens cDNA clone 647923 5' similar to SW YYAF\_BACSU P37518 HYPOTHETICAL 40.1 KD GTP-BINDING PROTEIN IN RPSF-SPO0J INTERGENIC REGION; mRNA sequence),

AA237053 (zs01c01.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 683904 5' similar to SW YBN5\_YEAST P38219 HYPOTHETICAL 44.2 KD PROTEIN IN SCO2-MRF1 INTERGENIC REGION), AA775776 (ad14e03.s1 Soares NbHFB Homo sapiens cDNA clone 878236 3' similar to TR P91917 P91917 W08E3.3; mRNA sequence), AL021878 (Homo sapiens DNA sequence from PAC 257I20 on chromosome 22q13.1-13.2; contains cytochrome P450 pseudogenes CYP2D7P, CYP2D8P, CYP2D6(D), TCF20, NADH ubiquinone oxidoreductase B14 subunit, ESTs, CA repeat, STS, GSS), and N32932 (yy10a02.s1 Homo sapiens cDNA clone 270794 3' similar to SW:YBN5\_YEAST P38219 HYPOTHETICAL 44.2 KD PROTEIN IN SCO2-MRF1 INTERGENIC REGION). The predicted amino acid sequence disclosed herein for qf122\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qf122\_3 protein demonstrated at least some similarity to sequences identified as W48670 (*Staphylococcus aureus* gbpA protein), Z92773 (W08E3.3 [*Caenorhabditis elegans*]), and Z92773 (predicted using Genefinder; Similarity to Yeast hypothetical 44.2 KD protein, putative GTP-binding protein (SW P38219); cDNA EST EMBL D64516 comes from this gene). Based upon sequence similarity, qf122\_3 proteins and each similar protein or peptide may share at least some activity. Analysis of protein motifs in SEQ ID NO:154 predicts an ATP/GTP-binding site motif A (P-loop) around amino acid 29 of SEQ ID NO:154.

20

#### Clone "qv538\_1"

A polynucleotide of the present invention has been identified as clone "qv538\_1". qv538\_1 was isolated from a human adult testes (embryonal carcinoma NT2D1 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qv538\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qv538\_1 protein").

The nucleotide sequence of qv538\_1 as presently determined is reported in SEQ ID NO:155, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qv538\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:156. Amino acids 8 to 20 of SEQ ID NO:156 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the

hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qv538\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
5 qv538\_1 should be approximately 2600 bp.

The nucleotide sequence disclosed herein for qv538\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qv538\_1 demonstrated at least some similarity with sequences identified as W44974 (zc22e11.r1 Soares senescent fibroblasts NbHSF Homo sapiens  
10 cDNA clone 323084 5' similar to SW:FKB2\_YEAST P32472 FK506-BINDING PROTEIN PRECURSOR; mRNA sequence), and Z62799 (H.sapiens CpG island DNA genomic MseI fragment, clone 73c8, reverse read cpg73c8.rt1a). The predicted amino acid sequence disclosed herein for qv538\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qv538\_1 protein  
15 demonstrated at least some similarity to sequences identified as AF04025 (FK506-binding protein [Mus musculus]) and W88556 (Secreted protein encoded by gene 23 clone HSQEO84). FK506-binding protein (or "FKBP") is the major high-affinity binding protein, in vertebrates, for the immunosuppressive drug FK506 (used to aid in organ transplantation acceptance among other indications). It exhibits peptidyl-prolyl cis-trans  
20 isomerase activity (PPIase or rotamase). PPIase is an enzyme that accelerates protein folding by catalyzing the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. Based upon sequence similarity, qv538\_1 proteins and each similar protein or peptide may share at least some activity. Analysis of protein motifs in SEQ ID NO:156 detects an endoplasmic reticulum targeting sequence around amino acid 208. Hidden  
25 Markov Model analysis detects an EF-hand calcium-binding domain at amino acids 183 to 211 of SEQ ID NO:156 (also found by motif analysis) and a FKBP-type peptidyl-prolyl cis-trans isomerase signatures/profile at amino acids 38 to 132 of SEQ ID NO:156. The nucleotide sequence of qv538\_1 indicates that it may contain an Alu repetitive element.

qv538\_1 protein was expressed in a COS cell expression system, and an expressed  
30 protein band of approximately 24 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "ys20\_1"

A polynucleotide of the present invention has been identified as clone "ys20\_1". ys20\_1 was isolated from a human adult thymus cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ys20\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ys20\_1 protein").

The nucleotide sequence of ys20\_1 as presently determined is reported in SEQ ID NO:157, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ys20\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:158. Amino acids 41 to 53 of SEQ ID NO:158 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 54. Amino acids 121 to 133 of SEQ ID NO:158 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning in that case at amino acid 134. Due to the hydrophobic nature of each of these predicted leader/signal sequences, each predicted leader/signal sequence is likely to act as a transmembrane domain should it not be separated from the remainder of the ys20\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ys20\_1 should be approximately 2229 bp.

The nucleotide sequence disclosed herein for ys20\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ys20\_1 demonstrated at least some similarity with sequences identified as B76357 (RPC111-15B19.TV RPC111 Homo sapiens genomic clone R-15B19, genomic survey sequence). Based upon sequence similarity, ys20\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the ys20\_1 protein sequence, centered around amino acid 205 of SEQ ID NO:158. The nucleotide sequence of ys20\_1 indicates that it may contain one or more mammalian transposon-like long terminal repeat elements, such as MCT1b/c.

Clone "as180\_1"

A polynucleotide of the present invention has been identified as clone "as180\_1". as180\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was

identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. as180\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "as180\_1 protein").

5       The nucleotide sequence of as180\_1 as presently determined is reported in SEQ ID NO:159. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the as180\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:160. Amino acids 168 to 180 of SEQ ID NO:160 are a predicted leader/signal sequence, with the predicted mature amino acid  
10       sequence beginning at amino acid 181. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the as180\_1 protein.

      The EcoRI/NotI restriction fragment obtainable from the deposit containing clone as180\_1 should be approximately 3580 bp.

15       The nucleotide sequence disclosed herein for as180\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. as180\_1 demonstrated at least some similarity with sequences identified as AB018279 (Homo sapiens mRNA for KIAA0736 protein, complete cds), S47919 (p87 = transporter-like protein [cattle, mRNA]), V89585 (EST clone CR618), and  
20       W28902 (53d11 Human retina cDNA randomly primed sublibrary Homo sapiens cDNA, mRNA sequence). The predicted amino acid sequence disclosed herein for as180\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted as180\_1 protein demonstrated at least some similarity to sequences identified as AB018279 (KIAA0736 protein [Homo sapiens]),  
25       L05435 (synaptic vesicle protein 2 [Rattus norvegicus]), S47919 (p87 [Bos sp.]), and W64538 (Human liver cell clone HP01293 protein). Synaptic vesicle protein 2 (SV2) is a membrane glycoprotein specifically localized to secretory vesicles in neurons and endocrine cells (Bajjalieh, S.M. *et al.*, 1992, *Science* Aug 28; 257(5074):1271-1273, which is incorporated by reference herein). Based upon sequence similarity, as180\_1 proteins and  
30       each similar protein or peptide may share at least some activity. Analysis of amino acid motifs detected a sugar-transport protein signature around amino acid 264 of SEQ ID NO:160, and hidden Markov Model analysis detected a sugar-transporter amino acid profile from amino acid 153 to amino acid 741 of SEQ ID NO:160. The TopPredII



computer program predicts twelve potential transmembrane domains within the as180\_1 protein sequence, centered around amino acids 181, 205, 248, 270, 308, 344, 432, 458, 605, 638, 654, and 710 of SEQ ID NO:160, respectively.

5        Deposit of Clones

Clones co62\_12, lo311\_8, ns197\_1, pj193\_5, pj317\_2, pt332\_1, qc297\_15, qg596\_12, and rb649\_3 were deposited on July 29, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 98825, from  
10 which each clone comprising a particular polynucleotide is obtainable.

Clones ca106\_19xx, ci52\_2, md124\_16, pk366\_7, pl741\_5, pp314\_19, pv35\_1, pw337\_6, rd610\_1, and rd810\_6 were deposited on August 11, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession  
15 number ATCC 98835, from which each clone comprising a particular polynucleotide is obtainable.

Clones cf85\_1, dd504\_18, np26\_3, pm412\_12, pm421\_3, pv6\_1, qs14\_3, qy338\_9, rc58\_1, and rd232\_5 were deposited on August 27, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an  
20 original deposit under the Budapest Treaty and were given the accession number ATCC 98850, from which each clone comprising a particular polynucleotide is obtainable.

Clones ck213\_12, pg195\_1, pw460\_5, qa136\_1, qy1261\_2, and rd432\_4 were deposited on October 8, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under  
25 the Budapest Treaty and were given the accession number ATCC 98918, from which each clone comprising a particular polynucleotide is obtainable.

Clones rb789\_14, yd137\_1, yd218\_1, ye11\_1, ye72\_1, ye78\_1, ye90\_1, yi62\_1, yk78\_1, yk251\_1, and yt14\_1 were deposited on December 15, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209  
30 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 207004, from which each clone comprising a particular polynucleotide is obtainable.

Clones bf157\_16, bk343\_2, cd205\_2, cw1292\_8, cw1475\_2, dd428\_4, dh1073\_12, dw78\_1, fh116\_11, fy356\_14, and iw66\_1 were deposited on February 4, 1999 with the

American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 207088, from which each clone comprising a particular polynucleotide is obtainable.

5 Clones kh13\_4, ko258\_4, kv10\_8, LL89\_3, mc300\_1, ml227\_1, mm367\_6, mt124\_3, nf56\_3, qy442\_2, rj214\_1, and rk80\_3 were deposited on February 4, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 207089, from which each clone comprising a particular  
10 polynucleotide is obtainable.

Clones au36\_42, bo549\_13, da529\_3, dm365\_3, fa171\_1, lp572\_2, pe246\_1, qf122\_3, qv538\_1, and ys20\_1 were deposited on April 2, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC  
15 207187, from which each clone comprising a particular polynucleotide is obtainable.

Clone as180\_1 was deposited on August 11, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC  
20 XXXXXX, from which the as180\_1 clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R. § 1.806.

25 Each clone has been transfected into separate bacterial cells (*E. coli*) in the above composite deposits. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or pNOTs vector depicted in Figures 1A and 1B, respectively, or in the case of clone qs14\_3,  
30 in the pCMVSPORT2 vector (Life Technologies, Inc., Rockville, MD 20850, U.S.A.) depicted in Figure 2. The pED6dpc2 vector ("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and

insertion of the M13 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences provided herein, or from a combination of those sequences. The sequence of an oligonucleotide probe that was used to isolate or to sequence each full-length clone is identified below, and should be most reliable in isolating the clone of interest.

<u>Clone</u>	<u>Probe Sequence</u>
co62_12	SEQ ID NO:161
lo311_8	SEQ ID NO:162
ns197_1	SEQ ID NO:163
pj193_5	SEQ ID NO:164
pj317_2	SEQ ID NO:165
pt332_1	SEQ ID NO:166
qc297_15	SEQ ID NO:167
qg596_12	SEQ ID NO:168
rb649_3	SEQ ID NO:169
ca106_19x	SEQ ID NO:170
ci52_2	SEQ ID NO:171
md124_16	SEQ ID NO:172
pk366_7	SEQ ID NO:173
pl741_5	SEQ ID NO:174
pp314_19	SEQ ID NO:175
pv35_1	SEQ ID NO:176
pw337_6	SEQ ID NO:177
rd610_1	SEQ ID NO:178
rd810_6	SEQ ID NO:179

	cf85_1	SEQ ID NO:180
	dd504_18	SEQ ID NO:181
	np26_3	SEQ ID NO:182
	pm412_12	SEQ ID NO:183
5	pm421_3	SEQ ID NO:184
	pv6_1	SEQ ID NO:185
	qs14_3	SEQ ID NO:186
	qy338_9	SEQ ID NO:187
	rc58_1	SEQ ID NO:188
10	rd232_5	SEQ ID NO:189
	ck213_12	SEQ ID NO:190
	pg195_1	SEQ ID NO:191
	pw460_5	SEQ ID NO:192
	qa136_1	SEQ ID NO:193
15	qy1261_2	SEQ ID NO:194
	rd432_4	SEQ ID NO:195
	rb789_14	SEQ ID NO:196
	yd137_1	SEQ ID NO:197
	ye11_1	SEQ ID NO:198
20	ye72_1	SEQ ID NO:199
	ye78_1	SEQ ID NO:200
	ye90_1	SEQ ID NO:201
	yk251_1	SEQ ID NO:202
	yt14_1	SEQ ID NO:203
25	bf157_16	SEQ ID NO:204
	bk343_2	SEQ ID NO:205
	cd205_2	SEQ ID NO:206
	cw1292_8	SEQ ID NO:207
	cw1475_2	SEQ ID NO:208
30	dd428_4	SEQ ID NO:209
	dh1073_12	SEQ ID NO:210
	dw78_1	SEQ ID NO:211
	fh116_11	SEQ ID NO:212
	fy356_14	SEQ ID NO:213

	iw66_1	SEQ ID NO:214
	kh13_4	SEQ ID NO:215
	ko258_4	SEQ ID NO:216
	kv10_8	SEQ ID NO:217
5	LL89_3	SEQ ID NO:218
	mc300_1	SEQ ID NO:219
	ml227_1	SEQ ID NO:220
	mm367_6	SEQ ID NO:221
	mt124_3	SEQ ID NO:222
10	nf56_3	SEQ ID NO:223
	qy442_2	SEQ ID NO:224
	rj214_14	SEQ ID NO:225
	rk80_3	SEQ ID NO:226
	au36_42	SEQ ID NO:227
15	bo549_13	SEQ ID NO:228
	da529_3	SEQ ID NO:229
	dm365_3	SEQ ID NO:230
	fa171_1	SEQ ID NO:231
	lp572_2	SEQ ID NO:232
20	pe246_1	SEQ ID NO:233
	qf122_3	SEQ ID NO:234
	qv538_1	SEQ ID NO:235
	ys20_1	SEQ ID NO:236
	as180_1	SEQ ID NO:237

25

In the sequences listed above which include an N at position 2, that position is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

30

The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;

- (b) It should be designed to have a  $T_m$  of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).

The oligonucleotide should preferably be labeled with  $\gamma$ -<sup>32</sup>P ATP (specific activity 6000 Ci/mmol) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a scintillation counter. Preferably, specific activity of the resulting probe should be approximately 4e+6 dpm/pmol.

10 The bacterial culture containing the pool of full-length clones should preferably be thawed and 100  $\mu$ l of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100  $\mu$ g/ml. The culture should preferably be grown to saturation at 37°C, and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions should preferably be plated to determine the  
15 dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100  $\mu$ g/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at 37°C. Other known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the  
20 colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100  $\mu$ g/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at  
25 a concentration greater than or equal to 1e+6 dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The  
30 filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein may be obtained by expression of the disclosed full-length polynucleotide (preferably those deposited with ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that

has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

The chromosomal location corresponding to the polynucleotide sequences disclosed herein may also be determined, for example by hybridizing appropriately  
5 labeled polynucleotides of the present invention to chromosomes *in situ*. It may also be possible to determine the corresponding chromosomal location for a disclosed polynucleotide by identifying significantly similar nucleotide sequences in public  
10 databases, such as expressed sequence tags (ESTs), that have already been mapped to particular chromosomal locations. For at least some of the polynucleotide sequences disclosed herein, public database sequences having at least some similarity to the  
15 polynucleotide of the present invention have been listed by database accession number. Searches using the GenBank accession numbers of these public database sequences can then be performed at an Internet site provided by the National Center for Biotechnology Information having the address <http://www.ncbi.nlm.nih.gov/UniGene/>, in order to  
20 identify "UniGene clusters" of overlapping sequences. Many of the "UniGene clusters" so identified will already have been mapped to particular chromosomal sites.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense  
25 polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, *Trends Pharmacol. Sci.* 15(7): 250-254; Lavarosky *et al.*, 1997, *Biochem. Mol. Med.* 62(1): 11-22; and Hampel, 1998, *Prog. Nucleic Acid Res. Mol. Biol.* 58: 1-39; all of which are incorporated by reference herein). The desired change in gene expression can also be achieved through the use of double-stranded ribonucleotide  
30 molecules having some complementarity to the mRNA transcribed from the gene, and which interfere with the transcription, stability, or expression of the mRNA ("RNA interference" or "RNAi"; Fire *et al.*, 1998, *Nature* 391 (6669): 806-811; Montgomery *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95 (26): 15502-15507; and Sharp, 1999, *Genes Dev.* 13 (2): 139-141; all of which are incorporated by reference herein). Transgenic animals that have  
multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are



also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through  
5 deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, *Bioessays* 14(9): 629-633; Zwaal *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 90(16): 7431-7435; Clark *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination,  
10 preferably detected by positive/negative genetic selection strategies (Mansour *et al.*, 1988, *Nature* 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for  
15 the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part  
20 or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of  
25 transmembrane domains in an amino acid sequence, domains which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most  
30 preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are

proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

5 In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* 266: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of*  
10 *Molecular Biology* 215: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* 3: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX  
15 platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely  
20 used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length  
25 one is  $Q=9$  for proteins and BLASTP, and  $Q=10$  for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is  $R=2$  for proteins and BLASTP, and  $R=10$  for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six,  
30 seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps.

The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

Species homologues of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuánez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90%

identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein and  
5 screening a suitable nucleic acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides that hybridize under reduced stringency conditions, more preferably stringent conditions, and most preferably highly  
10 stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) <sup>†</sup>	Hybridization Temperature and Buffer <sup>‡</sup>	Wash Temperature and Buffer <sup>‡</sup>
5	A	≥ 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
	B	<50	T <sub>B</sub> <sup>*</sup> ; 1xSSC	T <sub>B</sub> <sup>*</sup> ; 1xSSC
	C	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
	D	<50	T <sub>D</sub> <sup>*</sup> ; 1xSSC	T <sub>D</sub> <sup>*</sup> ; 1xSSC
	E	≥ 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
10	F	<50	T <sub>F</sub> <sup>*</sup> ; 1xSSC	T <sub>F</sub> <sup>*</sup> ; 1xSSC
	G	≥ 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
	H	<50	T <sub>H</sub> <sup>*</sup> ; 4xSSC	T <sub>H</sub> <sup>*</sup> ; 4xSSC
	I	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
	J	<50	T <sub>J</sub> <sup>*</sup> ; 4xSSC	T <sub>J</sub> <sup>*</sup> ; 4xSSC
15	K	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
	L	<50	T <sub>L</sub> <sup>*</sup> ; 2xSSC	T <sub>L</sub> <sup>*</sup> ; 2xSSC
	M	≥ 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
	N	<50	T <sub>N</sub> <sup>*</sup> ; 6xSSC	T <sub>N</sub> <sup>*</sup> ; 6xSSC
	O	≥ 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
20	P	<50	T <sub>P</sub> <sup>*</sup> ; 6xSSC	T <sub>P</sub> <sup>*</sup> ; 6xSSC
	Q	≥ 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
	R	<50	T <sub>R</sub> <sup>*</sup> ; 4xSSC	T <sub>R</sub> <sup>*</sup> ; 4xSSC

<sup>†</sup>: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

<sup>‡</sup>: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub>, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

<sup>\*</sup>T<sub>B</sub> - T<sub>R</sub>: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T<sub>m</sub>) of the hybrid, where T<sub>m</sub> is determined according to the following equations. For hybrids less than 18 base pairs in length, T<sub>m</sub>(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T<sub>m</sub>(°C) = 81.5 + 16.6(log<sub>10</sub>[Na<sup>+</sup>]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na<sup>+</sup>] is the concentration of sodium ions in the hybridization buffer ([Na<sup>+</sup>] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

The isolated polynucleotide encoding the protein of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* **19**, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* **185**, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial

strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from the Eastman Kodak Company (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to  
5 provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which  
10 are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by  
15 virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement,  
20 insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement,  
25 insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art



given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

### USES AND BIOLOGICAL ACTIVITY

5           The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies  
10       or vectors suitable for introduction of DNA).

#### Research Uses and Utilities

          The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express  
15       recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare  
20       with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for  
25       examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, those  
30       described in Gyuris *et al.*, 1993, *Cell* 75: 791-803 and in Rossi *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* 94: 8405-8410, all of which are incorporated by reference herein) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine  
5 levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the  
10 protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent  
15 grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to  
20 Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein  
25 or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention  
30 can be added to the medium in or on which the microorganism is cultured.

#### Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may

induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

- 10        Assays for T-cell or thymocyte proliferation include without limitation those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., *J. Immunol.* 137:3494-3500, 1986;
- 15        Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

- 20        Assays for cytokine production and /or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: *Polyclonal T cell stimulation*, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and *Measurement of mouse and human Interferon  $\gamma$* , Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

- 25        Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: *Measurement of Human and Murine Interleukin 2 and Interleukin 4*, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., *J. Exp. Med.* 173:1205-1211, 1991; Moreau et al., *Nature* 336:690-692, 1988; Greenberger et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983;
- 30        *Measurement of mouse and human interleukin 6* - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; *Measurement of human Interleukin 11* - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991;

Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

- Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

15      Immune Stimulating or Suppressing Activity

- A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

- 30      Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitis, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease.

Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present  
5 invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by  
10 suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and  
15 persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), *e.g.*, preventing  
20 high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys  
25 the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the  
30 molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term

tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

- 5       The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as
- 10       described in Lenschow *et al.*, Science 257:789-792 (1992) and Turka *et al.*, Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.
- 15       Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms.
- 20       Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from
- 25       the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/*lpr/lpr* mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and
- 30       murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune

response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain protein and  $\beta_2$

microglobulin protein or an MHC class II  $\alpha$  chain protein and an MHC class II  $\beta$  chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated  
 5 immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated  
 10 immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without  
 15 limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al.,  
 20 *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512, 1988; Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al., *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bowman et al., *J. Virology* 61:1992-1998; Takai et al., *J. Immunol.* 140:508-512, 1988; Bertagnoli et al.,  
 25 *Cellular Immunology* 133:327-341, 1991; Brown et al., *J. Immunol.* 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, *J. Immunol.* 144:3028-3033, 1990; and Assays for B cell function: *In vitro*  
 30 antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek,



D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

- 5 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995;
- 10 Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

- Assays for lymphocyte survival/apoptosis (which will identify, among others,
- 15 proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993;
- 20 Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

- Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

25

#### Hematopoiesis Regulating Activity

- A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell
- 30 lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid

cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and

Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

5

#### Tissue Growth Activity

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns,  
10 incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as  
15 well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal  
20 disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue  
25 destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in  
30 circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and

in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation

of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium ).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

#### Activin/Inhibin Activity

A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

#### Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. *J. Clin. Invest.* 95:1370-1376, 1995; Lind et al.

APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

#### Hemostatic and Thrombolytic Activity

5 A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting  
10 formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

15 Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### 20 Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands,  
25 receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant  
30 receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

10 Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

25

Cadherin/Tumor Invasion Suppressor Activity

Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus vulgaris and pemphigus foliaceus (auto-immune blistering skin diseases), Crohn's disease, and some developmental abnormalities.

30

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved



extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The cadherin domains bind calcium to form their tertiary structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this  
5 recognition site can change the specificity of a cadherin so that instead of recognizing only itself, the mutant molecule can now also bind to a different cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells  
10 become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal, restoring cells' adhesiveness to each other and to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas  
15 to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed  
20 in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different tissue in the body. Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the  
25 inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

Additionally, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block  
30 the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.

Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and polynucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable effects. Additionally, fragments of proteins of the present invention with cadherin activity, preferably truncated soluble cadherin fragments which have been found to be stable in the circulation of cancer patients, and polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al. Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

#### Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

#### Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s);

effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

#### ADMINISTRATION AND DOSING

A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical

compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in  
5 combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If  
10 administered sequentially, the attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical  
15 composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is  
20 administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention.  
25 When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid  
30 form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present

invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should  
5 contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

10 The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician  
15 will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu$ g to about 100  
20 mg (preferably about 0.1mg to about 10 mg, more preferably about 0.1  $\mu$ g to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is  
25 contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain  
30 polyclonal and monoclonal antibodies which specifically react with the protein. As used herein, the term "antibody" includes without limitation a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a single-chain antibody, a CDR-grafted antibody, a humanized antibody, or fragments thereof which bind to the indicated protein.

Such term also includes any other species derived from an antibody or antibody sequence which is capable of binding the indicated protein.

Antibodies to a particular protein can be produced by methods well known to those skilled in the art. For example, monoclonal antibodies can be produced by generation of  
5 antibody-producing hybridomas in accordance with known methods (see for example, Goding, 1983, *Monoclonal antibodies: principles and practice*, Academic Press Inc., New York; and Yokoyama, 1992, "Production of Monoclonal Antibodies" in *Current Protocols in Immunology*, Unit 2.5, Greene Publishing Assoc. and John Wiley & Sons). Polyclonal sera and antibodies can be produced by inoculation of a mammalian subject with the  
10 relevant protein or fragments thereof in accordance with known methods. Fragments of antibodies, receptors, or other reactive peptides can be produced from the corresponding antibodies by cleavage of and collection of the desired fragments in accordance with known methods (see for example, Goding, *supra*; and Andrew et al., 1992, "Fragmentation of Immunoglobulins" in *Current Protocols in Immunology*, Unit 2.8, Greene Publishing  
15 Assoc. and John Wiley & Sons). Chimeric antibodies and single chain antibodies can also be produced in accordance with known recombinant methods (see for example, 5,169,939, 5,194,594, and 5,576,184). Humanized antibodies can also be made from corresponding murine antibodies in accordance with well known methods (see for example, U.S. Patent Nos. 5,530,101, 5,585,089, and 5,693,762). Additionally, human antibodies may be  
20 produced in non-human animals such as mice that have been genetically altered to express human antibody molecules (see for example Fishwild *et al.*, 1996, *Nature Biotechnology* 14: 845-851; Mendez *et al.*, 1997, *Nature Genetics* 15: 146-156 (erratum *Nature Genetics* 16: 410); and U.S. Patents 5,877,397 and 5,625,126). Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide  
25 immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, *FEBS Lett.* 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful  
30 diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where

abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

5 For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably  
10 be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the  
15 methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical  
20 applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium  
25 sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other  
30 ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.



Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions  
5 from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of  
10 carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to  
15 provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in  
20 question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to  
25 humans, are desired patients for such treatment with proteins of the present invention.

The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of  
30 a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect

the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

5 Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

10 Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

Patent and literature references cited herein are incorporated by reference as if fully set forth.

15

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:1;
  - 5 (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 87 to nucleotide 821;
  - (c) the nucleotide sequence of SEQ ID NO:1 from nucleotide 120 to nucleotide 821;
  - (d) the nucleotide sequence of SEQ ID NO:1 from nucleotide 1 to  
10 nucleotide 1625;
  - (e) the nucleotide sequence of the full-length protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825;
  - (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;
  - 15 (g) the nucleotide sequence of a mature protein coding sequence of clone co62\_12 deposited under accession number ATCC 98825;
  - (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;
  - (i) a nucleotide sequence encoding a protein comprising the amino  
20 acid sequence of SEQ ID NO:2;
  - (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
  - (k) the nucleotide sequence of a polynucleotide that hybridizes under  
25 conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
  - (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and  
30 that has a length that is at least 25% of the length of SEQ ID NO:1.
2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.

3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:
  - (a) growing a culture of a host cell in a suitable culture medium, wherein the host cell has been transformed with the polynucleotide of claim 2; and
  - (b) purifying said protein from the culture.
6. A protein produced according to the process of claim 5.
7. An isolated polynucleotide encoding the protein of claim 6.
8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825.
9. A protein comprising an amino acid sequence selected from the group consisting of:
  - (a) the amino acid sequence of SEQ ID NO:2;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone co62\_12 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.
10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.
11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.
12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:3;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 9 to nucleotide 1013;
- (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 96 to nucleotide 1013;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- (f) the nucleotide sequence of a mature protein coding sequence of clone lo311\_8 deposited under accession number ATCC 98825;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
- (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
- (c) the amino acid sequence encoded by the cDNA insert of clone lo311\_8 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:5;
- 5 (b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 352 to nucleotide 825;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ns197\_1 deposited under accession number ATCC 98825;
- (d) a nucleotide sequence encoding the full-length protein encoded by  
10 the cDNA insert of clone ns197\_1 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
15 of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group  
25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:6, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:6; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
30 ns197\_1 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:7;
- (b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 86 to nucleotide 829;
- (c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 149 to nucleotide 829;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pj193\_5 deposited under accession number ATCC 98825;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:7.

17. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pj193\_5 deposited under accession number ATCC 98825;

the protein being substantially free from other mammalian proteins.

18. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:9;
- 5 (b) the nucleotide sequence of SEQ ID NO:9 from nucleotide 174 to nucleotide 1292;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pj317\_2 deposited under accession number ATCC 98825;
- (d) a nucleotide sequence encoding the full-length protein encoded by  
10 the cDNA insert of clone pj317\_2 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
15 of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under  
20 conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:9.

19. A protein comprising an amino acid sequence selected from the group  
25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:10, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
30 pj317\_2 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:



- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 7 to nucleotide 2517;
- (c) the nucleotide sequence of SEQ ID NO:11 from nucleotide 904 to nucleotide 2517;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pt332\_1 deposited under accession number ATCC 98825;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
- (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pt332\_1 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:13;
- 5 (b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 18 to nucleotide 257;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone qc297\_15 deposited under accession number ATCC 98825;
- (d) a nucleotide sequence encoding the full-length protein encoded by  
10 the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
15 of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:13.

25 23. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:14, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
  - 30 (c) the amino acid sequence encoded by the cDNA insert of clone qc297\_15 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:15;
- (b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 21 to nucleotide 2432;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone qg596\_12 deposited under accession number ATCC 98825;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone qg596\_12 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:17;
- (b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 339 to nucleotide 2105;
- (c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 501 to nucleotide 2105;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (f) the nucleotide sequence of a mature protein coding sequence of clone rb649\_3 deposited under accession number ATCC 98825;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
- (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
- (c) the amino acid sequence encoded by the cDNA insert of clone rb649\_3 deposited under accession number ATCC 98825;
- the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:19;
- 5 (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 509 to nucleotide 2467;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ca106\_19x deposited under accession number ATCC 98835;
- 10 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- 15 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:19.

25 29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - 30 (c) the amino acid sequence encoded by the cDNA insert of clone ca106\_19x deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 179 to nucleotide 802;
- (c) the nucleotide sequence of SEQ ID NO:21 from nucleotide 242 to nucleotide 802;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ci52\_2 deposited under accession number ATCC 98835;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
- (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and

(c) the amino acid sequence encoded by the cDNA insert of clone ci52\_2 deposited under accession number ATCC 98835; the protein being substantially free from other mammalian proteins.

5           32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 46 to nucleotide 714;
- 10           (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 538 to nucleotide 714;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding the full-length protein encoded by  
15           the cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (f) the nucleotide sequence of a mature protein coding sequence of clone md124\_16 deposited under accession number ATCC 98835;
- (g) a nucleotide sequence encoding a mature protein encoded by the  
20           cDNA insert of clone md124\_16 deposited under accession number ATCC 98835;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
25           of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under  
30           conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:24;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:24, the  
5 fragment comprising eight contiguous amino acids of SEQ ID NO:24; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
md124\_16 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

10 34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:25;
- (b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 92 to  
nucleotide 1726;
- 15 (c) the nucleotide sequence of SEQ ID NO:25 from nucleotide 1211 to  
nucleotide 1726;
- (d) the nucleotide sequence of the full-length protein coding sequence  
of clone pk366\_7 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding the full-length protein encoded by  
20 the cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;
- (f) the nucleotide sequence of a mature protein coding sequence of  
clone pk366\_7 deposited under accession number ATCC 98835;
- (g) a nucleotide sequence encoding a mature protein encoded by the  
cDNA insert of clone pk366\_7 deposited under accession number ATCC 98835;
- 25 (h) a nucleotide sequence encoding a protein comprising the amino  
acid sequence of SEQ ID NO:26;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight  
contiguous amino acids of SEQ ID NO:26;
- 30 (j) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees



C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:26;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 pk366\_7 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) the nucleotide sequence of SEQ ID NO:27;
- (b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 16 to nucleotide 1788;
- (c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 61 to nucleotide 1788;
- 20 (d) the nucleotide sequence of the full-length protein coding sequence of clone pl741\_5 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;
- (f) the nucleotide sequence of a mature protein coding sequence of  
25 clone pl741\_5 deposited under accession number ATCC 98835;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pl741\_5 deposited under accession number ATCC 98835;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- 30 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
- (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and
- (c) the amino acid sequence encoded by the cDNA insert of clone p1741\_5 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:29;
- (b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 629 to nucleotide 2338;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pp314\_19 deposited under accession number ATCC 98835;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pp314\_19 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

5 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
15 pp314\_19 deposited under accession number ATCC 98835;  
the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 20 (a) the nucleotide sequence of SEQ ID NO:31;
- (b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 158 to nucleotide 1102;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pv35\_1 deposited under accession number ATCC 98835;
- 25 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pv35\_1 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
30 of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:31.

5

41. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;
- (b) a fragment of the amino acid sequence of SEQ ID NO:32, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
pv35\_1 deposited under accession number ATCC 98835;  
the protein being substantially free from other mammalian proteins.

15

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 413 to  
nucleotide 733;
- (c) the nucleotide sequence of the full-length protein coding sequence  
20 of clone pw337\_6 deposited under accession number ATCC 98835;
- (d) a nucleotide sequence encoding the full-length protein encoded by  
the cDNA insert of clone pw337\_6 deposited under accession number ATCC  
98835;
- (e) a nucleotide sequence encoding a protein comprising the amino  
25 acid sequence of SEQ ID NO:34;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight  
contiguous amino acids of SEQ ID NO:34;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under  
30 conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 pw337\_6 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) the nucleotide sequence of SEQ ID NO:35;
- (b) the nucleotide sequence of SEQ ID NO:35 from nucleotide 678 to nucleotide 938;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone rd610\_1 deposited under accession number ATCC 98835;
- 20 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rd610\_1 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
25 of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 30 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:35.

45. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:36, the  
5 fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
rd610\_1 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

10 46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO.;
- (b) the nucleotide sequence of SEQ ID NO: from nucleotide 75 to  
nucleotide 494;
- 15 (c) the nucleotide sequence of SEQ ID NO: from nucleotide 447 to  
nucleotide 494;
- (d) the nucleotide sequence of the full-length protein coding sequence  
of clone rd810\_6 deposited under accession number ATCC 98835;
- (e) a nucleotide sequence encoding the full-length protein encoded by  
20 the cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835;
- (f) the nucleotide sequence of a mature protein coding sequence of  
clone rd810\_6 deposited under accession number ATCC 98835;
- (g) a nucleotide sequence encoding a mature protein encoded by the  
cDNA insert of clone rd810\_6 deposited under accession number ATCC 98835;
- 25 (h) a nucleotide sequence encoding a protein comprising the amino  
acid sequence of SEQ ID NO.;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO., the fragment comprising eight  
contiguous amino acids of SEQ ID NO.;
- 30 (j) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:.

47. A protein comprising an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:, the fragment comprising eight contiguous amino acids of SEQ ID NO:; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 rd810\_6 deposited under accession number ATCC 98835;
- the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 181 to nucleotide 1080;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone cf85\_1 deposited under accession number ATCC 98850;
- 20 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
25 of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 30 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cf85\_1 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

10

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:41;
- (b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 161 to  
15 nucleotide 1348;
- (c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 599 to nucleotide 1348;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850;
- 20 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- (f) the nucleotide sequence of a mature protein coding sequence of clone dd504\_18 deposited under accession number ATCC 98850;
- 25 (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
30 of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and



(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

5

51. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:42, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dd504\_18 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

15

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:43;
- (b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 70 to  
nucleotide 1386;
- (c) the nucleotide sequence of the full-length protein coding sequence  
20 of clone np26\_3 deposited under accession number ATCC 98850;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone np26\_3 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding a protein comprising the amino  
25 acid sequence of SEQ ID NO:44;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under  
30 conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:44;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 np26\_3 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) the nucleotide sequence of SEQ ID NO:45;
- (b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 60 to nucleotide 3515;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pm412\_12 deposited under accession number ATCC 98850;
- 20 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- 25 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
30 C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm412\_12 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 1490 to nucleotide 1780;
- (c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 1556 to nucleotide 1780;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pm421\_3 deposited under accession number ATCC 98850;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

5

57. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm421\_3 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

15

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 64 to nucleotide 486;
- (c) the nucleotide sequence of SEQ ID NO:49 from nucleotide 217 to  
20 nucleotide 486;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding the full-length protein encoded by  
25 the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pv6\_1 deposited under accession number ATCC 98850;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pv6\_1 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a protein comprising the amino  
30 acid sequence of SEQ ID NO:50;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

5 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
15 pv6\_1 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 20 (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 379 to nucleotide 3783;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 460 to nucleotide 3783;
- 25 (d) the nucleotide sequence of SEQ ID NO:51 from nucleotide 1983 to nucleotide 3938;
- (e) the nucleotide sequence of the full-length protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850;
- (f) a nucleotide sequence encoding the full-length protein encoded by  
30 the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;
- (g) the nucleotide sequence of a mature protein coding sequence of clone qs14\_3 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;

- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:52;
- (b) the amino acid sequence of SEQ ID NO:52 from amino acid 536 to amino acid 1135;
- (c) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
- (d) the amino acid sequence encoded by the cDNA insert of clone qs14\_3 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:53;
- (b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 1 to nucleotide 843;
- (c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 469 to nucleotide 843;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qy338\_9 deposited under accession number ATCC 98850;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- (f) the nucleotide sequence of a mature protein coding sequence of clone qy338\_9 deposited under accession number ATCC 98850;
- 5 (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;
- 10 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 15 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.
- 20 63. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:54;
- (b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone qy338\_9 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.
64. An isolated polynucleotide comprising a nucleotide sequence selected from
- 30 the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:55;
- (b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 283 to nucleotide 906;

- (c) the nucleotide sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 906;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850;
- 5 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;
- (f) the nucleotide sequence of a mature protein coding sequence of clone rc58\_1 deposited under accession number ATCC 98850;
- 10 (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone rc58\_1 deposited under accession number ATCC 98850;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight
- 15 contiguous amino acids of SEQ ID NO:56;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 20 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:55.
65. A protein comprising an amino acid sequence selected from the group
- 25 consisting of:
- (a) the amino acid sequence of SEQ ID NO:56;
- (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
- (c) the amino acid sequence encoded by the cDNA insert of clone
- 30 rc58\_1 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:



- (a) the nucleotide sequence of SEQ ID NO:57;
- (b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 56 to nucleotide 973;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone rd232\_5 deposited under accession number ATCC 98850;
- 5 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- 10 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 15 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:57.
- 20
67. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:58;
- (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone rd232\_5 deposited under accession number ATCC 98850;
- the protein being substantially free from other mammalian proteins.
- 30
68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:59;
- (b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 893 to nucleotide 2596;

- (c) the nucleotide sequence of the full-length protein coding sequence of clone ck213\_12 deposited under accession number ATCC 98918;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:59.
69. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:60;
- (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ck213\_12 deposited under accession number ATCC 98918;
- the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 29 to nucleotide 1750;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pg195\_1 deposited under accession number ATCC 98918;

- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;
- 5 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 10 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:61.

15

71. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
- (b) a fragment of the amino acid sequence of SEQ ID NO:62, the
- 20 fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pg195\_1 deposited under accession number ATCC 98918;
- the protein being substantially free from other mammalian proteins.

25

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:63;
- (b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 1147 to nucleotide 1440;
- 30 (c) the nucleotide sequence of SEQ ID NO:63 from nucleotide 1234 to nucleotide 1440;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pw460\_5 deposited under accession number ATCC 98918;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:64;
- (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pw460\_5 deposited under accession number ATCC 98918;
- the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 46 to nucleotide 1356;

- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 127 to nucleotide 1356;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918;
- 5 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
- (f) the nucleotide sequence of a mature protein coding sequence of clone qa136\_1 deposited under accession number ATCC 98918;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
- 10 (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- 15 (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.
- 20

75. A protein comprising an amino acid sequence selected from the group consisting of:

25

- (a) the amino acid sequence of SEQ ID NO:66;
- (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qa136\_1 deposited under accession number ATCC 98918;
- 30 the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:67;
- (b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 206 to nucleotide 1624;
- (c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 542 to nucleotide 1624;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;
- (f) the nucleotide sequence of a mature protein coding sequence of clone qy1261\_2 deposited under accession number ATCC 98918;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
- (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qy1261\_2 deposited under accession number ATCC 98918;

the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 5 (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 1359 to nucleotide 1817;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone rd432\_4 deposited under accession number ATCC 98918;
- 10 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
15 of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:69.

25 79. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:70;
- (b) a fragment of the amino acid sequence of SEQ ID NO:70, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:70; and
- 30 (c) the amino acid sequence encoded by the cDNA insert of clone rd432\_4 deposited under accession number ATCC 98918;

the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:71;
- (b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 884 to nucleotide 1195;
- (c) the nucleotide sequence of SEQ ID NO:71 from nucleotide 947 to nucleotide 1195;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone rb789\_14 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004;
- (f) the nucleotide sequence of a mature protein coding sequence of clone rb789\_14 deposited under accession number ATCC 207004;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:72;
- (b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and



(c) the amino acid sequence encoded by the cDNA insert of clone rb789\_14 deposited under accession number ATCC 207004; the protein being substantially free from other mammalian proteins.

- 5           82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:73;
  - (b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 69 to nucleotide 374;
  - 10           (c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 186 to nucleotide 374;
  - (d) the nucleotide sequence of the full-length protein coding sequence of clone yd137\_1 deposited under accession number ATCC 207004;
  - (e) a nucleotide sequence encoding the full-length protein encoded by  
15           the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;
  - (f) the nucleotide sequence of a mature protein coding sequence of clone yd137\_1 deposited under accession number ATCC 207004;
  - (g) a nucleotide sequence encoding a mature protein encoded by the  
20           cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;
  - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
  - (i) a nucleotide sequence encoding a protein comprising a fragment  
25           of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
  - (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
  - (k) the nucleotide sequence of a polynucleotide that hybridizes under  
30           conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:74, the  
5 fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone yd137\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

10 84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 8 to  
nucleotide 343;
- 15 (c) the nucleotide sequence of SEQ ID NO:75 from nucleotide 50 to  
nucleotide 343;
- (d) the nucleotide sequence of the full-length protein coding sequence  
of clone yd218\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding the full-length protein encoded by  
20 the cDNA insert of clone yd218\_1 deposited under accession number ATCC  
207004;
- (f) the nucleotide sequence of a mature protein coding sequence of  
clone yd218\_1 deposited under accession number ATCC 207004;
- (g) a nucleotide sequence encoding a mature protein encoded by the  
25 cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
- (h) a nucleotide sequence encoding a protein comprising the amino  
acid sequence of SEQ ID NO:76;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight  
30 contiguous amino acids of SEQ ID NO:76;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:75.

5

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:76, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone yd218\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

15

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:77;
- (b) the nucleotide sequence of SEQ ID NO:77 from nucleotide 84 to  
nucleotide 1679;
- (c) the nucleotide sequence of the full-length protein coding sequence  
20 of clone ye11\_1 deposited under accession number ATCC 207004;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ye11\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding a protein comprising the amino  
25 acid sequence of SEQ ID NO:78;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under  
30 conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:77.

87. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:78;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ye11\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:79;
  - (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 72 to nucleotide 1646;
  - (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 180 to nucleotide 1646;
  - (d) the nucleotide sequence of the full-length protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004;
  - (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
  - (f) the nucleotide sequence of a mature protein coding sequence of clone ye72\_1 deposited under accession number ATCC 207004;
  - (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ye72\_1 deposited under accession number ATCC 207004;
  - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
  - (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

5 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:80;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
15 ye72\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

90. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 20 (a) the nucleotide sequence of SEQ ID NO:81;
- (b) the nucleotide sequence of SEQ ID NO:81 from nucleotide 954 to nucleotide 2423;
- (c) the nucleotide sequence of SEQ ID NO:81 from nucleotide 1224 to nucleotide 2423;
- 25 (d) the nucleotide sequence of the full-length protein coding sequence of clone ye78\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- (f) the nucleotide sequence of a mature protein coding sequence of  
30 clone ye78\_1 deposited under accession number ATCC 207004;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:82;

- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;
- 5 (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and
- 10 that has a length that is at least 25% of the length of SEQ ID NO:81.

91. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:82;
- 15 (b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ye78\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

20

92. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:83;
- (b) the nucleotide sequence of SEQ ID NO:83 from nucleotide 176 to
- 25 nucleotide 1321;
- (c) the nucleotide sequence of SEQ ID NO:83 from nucleotide 233 to nucleotide 1321;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004;
- 30 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ye90\_1 deposited under accession number ATCC 207004;

- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:84;
- 5 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees
- 10 C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:83.
- 15
93. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:84;
- (b) a fragment of the amino acid sequence of SEQ ID NO:84, the
- 20 fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ye90\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.
- 25
94. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:85;
- (b) the nucleotide sequence of SEQ ID NO:85 from nucleotide 105 to nucleotide 605;
- 30 (c) the nucleotide sequence of the full-length protein coding sequence of clone yi62\_1 deposited under accession number ATCC 207004;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;

- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:85.

95. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:86;
- (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yi62\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

96. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:87;
- (b) the nucleotide sequence of SEQ ID NO:87 from nucleotide 223 to nucleotide 798;
- (c) the nucleotide sequence of SEQ ID NO:87 from nucleotide 430 to nucleotide 798;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;



- (f) the nucleotide sequence of a mature protein coding sequence of clone yk78\_1 deposited under accession number ATCC 207004;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;
- 5 (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- 10 (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and
- 15 that has a length that is at least 25% of the length of SEQ ID NO:87.

97. A protein comprising an amino acid sequence selected from the group consisting of:
- 20 (a) the amino acid sequence of SEQ ID NO:88;
- (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yk78\_1 deposited under accession number ATCC 207004;
- 25 the protein being substantially free from other mammalian proteins.

98. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:89;
- 30 (b) the nucleotide sequence of SEQ ID NO:89 from nucleotide 211 to nucleotide 942;
- (c) the nucleotide sequence of SEQ ID NO:89 from nucleotide 298 to nucleotide 942;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
- (f) the nucleotide sequence of a mature protein coding sequence of clone yk251\_1 deposited under accession number ATCC 207004;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:90;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:89.

99. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:90;
- (b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and
- (c) the amino acid sequence encoded by the cDNA insert of clone yk251\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.

100. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:91;

- (b) the nucleotide sequence of SEQ ID NO:91 from nucleotide 149 to nucleotide 784;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone yt14\_1 deposited under accession number ATCC 207004;
- 5 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
10 of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 15 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:91.
- 20 101. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:92;
- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone yt14\_1 deposited under accession number ATCC 207004;
- the protein being substantially free from other mammalian proteins.
102. An isolated polynucleotide comprising a nucleotide sequence selected from  
30 the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:93;
- (b) the nucleotide sequence of SEQ ID NO:93 from nucleotide 89 to nucleotide 1441;

- (c) the nucleotide sequence of the full-length protein coding sequence of clone bf157\_16 deposited under accession number ATCC 207088;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:93.
103. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:94;
- (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bf157\_16 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

104. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:95;
- (b) the nucleotide sequence of SEQ ID NO:95 from nucleotide 219 to nucleotide 629;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone bk343\_2 deposited under accession number ATCC 207088;

- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;
- 5 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:96;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;
- 10 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and
- 15 that has a length that is at least 25% of the length of SEQ ID NO:95.

105. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:96;
- 20 (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bk343\_2 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

25

106. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:97;
- (b) the nucleotide sequence of SEQ ID NO:97 from nucleotide 556 to
- 30 nucleotide 951;
- (c) the nucleotide sequence of SEQ ID NO:97 from nucleotide 868 to nucleotide 951;
- (d) the nucleotide sequence of SEQ ID NO:97 from nucleotide 9 to nucleotide 1295;

- (e) the nucleotide sequence of the full-length protein coding sequence of clone cd205\_2 deposited under accession number ATCC 207088;
- (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
- (g) the nucleotide sequence of a mature protein coding sequence of clone cd205\_2 deposited under accession number ATCC 207088;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:98;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:97.
107. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:98;
- (b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cd205\_2 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

108. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:99;

- (b) the nucleotide sequence of SEQ ID NO:99 from nucleotide 216 to nucleotide 443;
- (c) the nucleotide sequence of SEQ ID NO:99 from nucleotide 306 to nucleotide 443;
- 5 (d) the nucleotide sequence of the full-length protein coding sequence of clone cw1292\_8 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
- 10 (f) the nucleotide sequence of a mature protein coding sequence of clone cw1292\_8 deposited under accession number ATCC 207088;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- 15 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 20 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and
- 25 that has a length that is at least 25% of the length of SEQ ID NO:99.

109. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
- 30 (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cw1292\_8 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

110. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:101;
- 5 (b) the nucleotide sequence of SEQ ID NO:101 from nucleotide 2136 to nucleotide 2447;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone cw1475\_2 deposited under accession number ATCC 207088;
- (d) a nucleotide sequence encoding the full-length protein encoded by  
10 the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (f) a nucleotide sequence encoding a protein comprising a fragment  
15 of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:101.

25 111. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:102, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
  - 30 (c) the amino acid sequence encoded by the cDNA insert of clone cw1475\_2 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.



112. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:103;
- (b) the nucleotide sequence of SEQ ID NO:103 from nucleotide 310 to nucleotide 954;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone dd428\_4 deposited under accession number ATCC 207088;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:103.

113. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dd428\_4 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

114. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:105;
- (b) the nucleotide sequence of SEQ ID NO:105 from nucleotide 1698 to nucleotide 1895;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone dh1073\_12 deposited under accession number ATCC 207088;
- 5 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- 10 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 15 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:105.
- 20

115. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:106;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dh1073\_12 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

30

116. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:107;

- (b) the nucleotide sequence of SEQ ID NO:107 from nucleotide 423 to nucleotide 791;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone dw78\_1 deposited under accession number ATCC 207088;
- 5 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:108;
- 10 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 15 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:107.
- 20
117. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:108;
- (b) a fragment of the amino acid sequence of SEQ ID NO:108, the
- 25 fragment comprising eight contiguous amino acids of SEQ ID NO:108; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dw78\_1 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.
- 30
118. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:109;
- (b) the nucleotide sequence of SEQ ID NO:109 from nucleotide 96 to nucleotide 944;

- (c) the nucleotide sequence of the full-length protein coding sequence of clone fh116\_11 deposited under accession number ATCC 207088;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:109.

119. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;
- (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fh116\_11 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

120. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:111;
- (b) the nucleotide sequence of SEQ ID NO:111 from nucleotide 150 to nucleotide 1610;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone fy356\_14 deposited under accession number ATCC 207088;

- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
- 5 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- 10 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and
- 15 that has a length that is at least 25% of the length of SEQ ID NO:111.

121. A protein comprising an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:112;
- (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fy356\_14 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

25

122. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:113;
- 30 (b) the nucleotide sequence of SEQ ID NO:113 from nucleotide 49 to nucleotide 669;
- (c) the nucleotide sequence of SEQ ID NO:113 from nucleotide 112 to nucleotide 669;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;
- (f) the nucleotide sequence of a mature protein coding sequence of clone iw66\_1 deposited under accession number ATCC 207088;
- 5 (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:114;
- (i) a nucleotide sequence encoding a protein comprising a fragment
- 10 of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 15 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:113.
- 20 123. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:114;
- (b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone iw66\_1 deposited under accession number ATCC 207088;
- the protein being substantially free from other mammalian proteins.

124. An isolated polynucleotide comprising a nucleotide sequence selected from
- 30 the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:115;
- (b) the nucleotide sequence of SEQ ID NO:115 from nucleotide 165 to nucleotide 416;

- (c) the nucleotide sequence of the full-length protein coding sequence of clone kh13\_4 deposited under accession number ATCC 207089;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
- 5 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;
- 10 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and
- 15 that has a length that is at least 25% of the length of SEQ ID NO:115.

125. A protein comprising an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:116;
- (b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and
- (c) the amino acid sequence encoded by the cDNA insert of clone kh13\_4 deposited under accession number ATCC 207089;
- 25 the protein being substantially free from other mammalian proteins.

126. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:117;
- 30 (b) the nucleotide sequence of SEQ ID NO:117 from nucleotide 204 to nucleotide 602;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ko258\_4 deposited under accession number ATCC 207089;

- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- 5 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;
- 10 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and
- 15 that has a length that is at least 25% of the length of SEQ ID NO:117.

127. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;
- 20 (b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ko258\_4 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.

25

128. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:119;
- (b) the nucleotide sequence of SEQ ID NO:119 from nucleotide 434 to
- 30 nucleotide 739;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone kv10\_8 deposited under accession number ATCC 207089;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;



- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:120;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120, the fragment comprising eight contiguous amino acids of SEQ ID NO:120;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:119.
129. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:120;
- (b) a fragment of the amino acid sequence of SEQ ID NO:120, the fragment comprising eight contiguous amino acids of SEQ ID NO:120; and
- (c) the amino acid sequence encoded by the cDNA insert of clone kv10\_8 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.
130. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:121;
- (b) the nucleotide sequence of SEQ ID NO:121 from nucleotide 149 to nucleotide 310;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone LL89\_3 deposited under accession number ATCC 207089;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:122;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122, the fragment comprising eight contiguous amino acids of SEQ ID NO:122;

5 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and  
10 that has a length that is at least 25% of the length of SEQ ID NO:121.

131. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:122;
- 15 (b) a fragment of the amino acid sequence of SEQ ID NO:122, the fragment comprising eight contiguous amino acids of SEQ ID NO:122; and
- (c) the amino acid sequence encoded by the cDNA insert of clone LL89\_3 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.

20

132. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:123;
- (b) the nucleotide sequence of SEQ ID NO:123 from nucleotide 22 to  
25 nucleotide 288;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone mc300\_1 deposited under accession number ATCC 207089;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC  
30 207089;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:124;

- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124, the fragment comprising eight contiguous amino acids of SEQ ID NO:124;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:123.

133. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:124;
- (b) a fragment of the amino acid sequence of SEQ ID NO:124, the fragment comprising eight contiguous amino acids of SEQ ID NO:124; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mc300\_1 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.

134. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:125;
- (b) the nucleotide sequence of SEQ ID NO:125 from nucleotide 200 to nucleotide 2449;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ml227\_1 deposited under accession number ATCC 207089;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:126;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126, the fragment comprising eight contiguous amino acids of SEQ ID NO:126;

5 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and  
10 that has a length that is at least 25% of the length of SEQ ID NO:125.

135. A protein comprising an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:126;  
(b) a fragment of the amino acid sequence of SEQ ID NO:126, the fragment comprising eight contiguous amino acids of SEQ ID NO:126; and  
(c) the amino acid sequence encoded by the cDNA insert of clone ml227\_1 deposited under accession number ATCC 207089;  
the protein being substantially free from other mammalian proteins.

20

136. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:127;  
(b) the nucleotide sequence of SEQ ID NO:127 from nucleotide 82 to  
25 nucleotide 1980;  
(c) the nucleotide sequence of the full-length protein coding sequence of clone mm367\_6 deposited under accession number ATCC 207089;  
(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC  
30 207089;  
(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:128;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128, the fragment comprising eight contiguous amino acids of SEQ ID NO:128;

5 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and  
10 that has a length that is at least 25% of the length of SEQ ID NO:127.

137. A protein comprising an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:128;  
(b) a fragment of the amino acid sequence of SEQ ID NO:128, the fragment comprising eight contiguous amino acids of SEQ ID NO:128; and  
(c) the amino acid sequence encoded by the cDNA insert of clone mm367\_6 deposited under accession number ATCC 207089;  
the protein being substantially free from other mammalian proteins.

20

138. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 25 (a) the nucleotide sequence of SEQ ID NO:129;  
(b) the nucleotide sequence of SEQ ID NO:129 from nucleotide 125 to nucleotide 856;  
(c) the nucleotide sequence of the full-length protein coding sequence of clone mt124\_3 deposited under accession number ATCC 207089;  
(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC  
30 207089;  
(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:130;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130;

5 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and  
10 that has a length that is at least 25% of the length of SEQ ID NO:129.

139. A protein comprising an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:130;
- (b) a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mt124\_3 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.

20

140. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:131;
- 25 (b) the nucleotide sequence of SEQ ID NO:131 from nucleotide 856 to nucleotide 2940;
- (c) the nucleotide sequence of SEQ ID NO:131 from nucleotide 901 to nucleotide 2940;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089;
- 30 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
- (f) the nucleotide sequence of a mature protein coding sequence of clone nf56\_3 deposited under accession number ATCC 207089;

- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:132;
- 5 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132, the fragment comprising eight contiguous amino acids of SEQ ID NO:132;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 10 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:131.
- 15 141. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:132;
- (b) a fragment of the amino acid sequence of SEQ ID NO:132, the
- 20 fragment comprising eight contiguous amino acids of SEQ ID NO:132; and
- (c) the amino acid sequence encoded by the cDNA insert of clone nf56\_3 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.
- 25 142. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:133;
- (b) the nucleotide sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 448;
- 30 (c) the nucleotide sequence of SEQ ID NO:133 from nucleotide 167 to nucleotide 448;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;

5 (f) the nucleotide sequence of a mature protein coding sequence of clone qy442\_2 deposited under accession number ATCC 207089;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:134;

10 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134, the fragment comprising eight contiguous amino acids of SEQ ID NO:134;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

15 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:133.

20

143. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:134;

25 (b) a fragment of the amino acid sequence of SEQ ID NO:134, the fragment comprising eight contiguous amino acids of SEQ ID NO:134; and

(c) the amino acid sequence encoded by the cDNA insert of clone qy442\_2 deposited under accession number ATCC 207089;

the protein being substantially free from other mammalian proteins.

30 144. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:135;

(b) the nucleotide sequence of SEQ ID NO:135 from nucleotide 28 to nucleotide 777;



- (c) the nucleotide sequence of SEQ ID NO:135 from nucleotide 73 to nucleotide 777;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089;
- 5 (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- (f) the nucleotide sequence of a mature protein coding sequence of clone rj214\_14 deposited under accession number ATCC 207089;
- 10 (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:136;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136, the fragment comprising eight contiguous amino acids of SEQ ID NO:136;
- 15 (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 20 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:135.
- 25 145. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:136;
- (b) a fragment of the amino acid sequence of SEQ ID NO:136, the fragment comprising eight contiguous amino acids of SEQ ID NO:136; and
- 30 (c) the amino acid sequence encoded by the cDNA insert of clone rj214\_14 deposited under accession number ATCC 207089;
- the protein being substantially free from other mammalian proteins.

146. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:137;
- (b) the nucleotide sequence of SEQ ID NO:137 from nucleotide 179 to  
5 nucleotide 745;
- (c) the nucleotide sequence of SEQ ID NO:137 from nucleotide 233 to  
nucleotide 745;
- (d) the nucleotide sequence of the full-length protein coding sequence  
of clone rk80\_3 deposited under accession number ATCC 207089;
- 10 (e) a nucleotide sequence encoding the full-length protein encoded by  
the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;
- (f) the nucleotide sequence of a mature protein coding sequence of  
clone rk80\_3 deposited under accession number ATCC 207089;
- (g) a nucleotide sequence encoding a mature protein encoded by the  
15 cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089;
- (h) a nucleotide sequence encoding a protein comprising the amino  
acid sequence of SEQ ID NO:138;
- (i) a nucleotide sequence encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO:138, the fragment comprising eight  
20 contiguous amino acids of SEQ ID NO:138;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under  
conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under  
25 conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees  
C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and  
that has a length that is at least 25% of the length of SEQ ID NO:137.

147. A protein comprising an amino acid sequence selected from the group  
30 consisting of:

- (a) the amino acid sequence of SEQ ID NO:138;
- (b) a fragment of the amino acid sequence of SEQ ID NO:138, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:138; and

(c) the amino acid sequence encoded by the cDNA insert of clone rk80\_3 deposited under accession number ATCC 207089; the protein being substantially free from other mammalian proteins.

5 148. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:139;
- (b) the nucleotide sequence of SEQ ID NO:139 from nucleotide 1017 to nucleotide 1274;
- 10 (c) the nucleotide sequence of the full-length protein coding sequence of clone au36\_42 deposited under accession number ATCC 207187;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;
- 15 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:140;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140, the fragment comprising eight contiguous amino acids of SEQ ID NO:140;
- 20 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and
- 25 that has a length that is at least 25% of the length of SEQ ID NO:139.

149. A protein comprising an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:140;
- (b) a fragment of the amino acid sequence of SEQ ID NO:140, the fragment comprising eight contiguous amino acids of SEQ ID NO:140; and
- (c) the amino acid sequence encoded by the cDNA insert of clone au36\_42 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins.

150. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 5 (a) the nucleotide sequence of SEQ ID NO:141;
- (b) the nucleotide sequence of SEQ ID NO:141 from nucleotide 580 to nucleotide 774;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone bo549\_13 deposited under accession number ATCC 207187;
- 10 (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:142;
- 15 (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142, the fragment comprising eight contiguous amino acids of SEQ ID NO:142;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 20 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:141.

25 151. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:142;
- (b) a fragment of the amino acid sequence of SEQ ID NO:142, the
- 30 fragment comprising eight contiguous amino acids of SEQ ID NO:142; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bo549\_13 deposited under accession number ATCC 207187;

the protein being substantially free from other mammalian proteins.

152. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:143;
- (b) the nucleotide sequence of SEQ ID NO:143 from nucleotide 172 to nucleotide 969;
- (c) the nucleotide sequence of SEQ ID NO:143 from nucleotide 385 to nucleotide 969;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187;
- (f) the nucleotide sequence of a mature protein coding sequence of clone da529\_3 deposited under accession number ATCC 207187;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:144;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144, the fragment comprising eight contiguous amino acids of SEQ ID NO:144;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:143.

153. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:144;
- (b) a fragment of the amino acid sequence of SEQ ID NO:144, the fragment comprising eight contiguous amino acids of SEQ ID NO:144; and

(c) the amino acid sequence encoded by the cDNA insert of clone da529\_3 deposited under accession number ATCC 207187; the protein being substantially free from other mammalian proteins.

- 5           154. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence of SEQ ID NO:145;
  - (b) the nucleotide sequence of SEQ ID NO:145 from nucleotide 329 to nucleotide 667;
  - 10           (c) the nucleotide sequence of SEQ ID NO:145 from nucleotide 368 to nucleotide 667;
  - (d) the nucleotide sequence of the full-length protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187;
  - (e) a nucleotide sequence encoding the full-length protein encoded by  
15           the cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;
  - (f) the nucleotide sequence of a mature protein coding sequence of clone dm365\_3 deposited under accession number ATCC 207187;
  - (g) a nucleotide sequence encoding a mature protein encoded by the  
20           cDNA insert of clone dm365\_3 deposited under accession number ATCC 207187;
  - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:146;
  - (i) a nucleotide sequence encoding a protein comprising a fragment  
25           of the amino acid sequence of SEQ ID NO:146, the fragment comprising eight contiguous amino acids of SEQ ID NO:146;
  - (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
  - (k) the nucleotide sequence of a polynucleotide that hybridizes under  
30           conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:145.

155. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:146;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:146, the  
5 fragment comprising eight contiguous amino acids of SEQ ID NO:146; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
dm365\_3 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins.

10 156. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:147;
- (b) the nucleotide sequence of SEQ ID NO:147 from nucleotide 103 to  
nucleotide 1368;

15 (c) the nucleotide sequence of the full-length protein coding sequence of clone fa171\_1 deposited under accession number ATCC 207187;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;

20 (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:148;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148, the fragment comprising eight contiguous amino acids of SEQ ID NO:148;

25 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

30 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:147.

157. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:148;
- (b) a fragment of the amino acid sequence of SEQ ID NO:148, the fragment comprising eight contiguous amino acids of SEQ ID NO:148; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fa171\_1 deposited under accession number ATCC 207187;
- 5 the protein being substantially free from other mammalian proteins.

158. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 10 (a) the nucleotide sequence of SEQ ID NO:149;
- (b) the nucleotide sequence of SEQ ID NO:149 from nucleotide 190 to nucleotide 1407;
- (c) the nucleotide sequence of SEQ ID NO:149 from nucleotide 463 to nucleotide 1407;
- 15 (d) the nucleotide sequence of the full-length protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
- 20 (f) the nucleotide sequence of a mature protein coding sequence of clone lp572\_2 deposited under accession number ATCC 207187;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone lp572\_2 deposited under accession number ATCC 207187;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:150;
- 25 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150, the fragment comprising eight contiguous amino acids of SEQ ID NO:150;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- 30 (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees



C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:149.

159. A protein comprising an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:150;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:150, the fragment comprising eight contiguous amino acids of SEQ ID NO:150; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 lp572\_2 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins.

160. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) the nucleotide sequence of SEQ ID NO:151;
- (b) the nucleotide sequence of SEQ ID NO:151 from nucleotide 301 to nucleotide 1035;
- (c) the nucleotide sequence of SEQ ID NO:151 from nucleotide 916 to nucleotide 1035;
- 20 (d) the nucleotide sequence of the full-length protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;
- 25 (f) the nucleotide sequence of a mature protein coding sequence of clone pe246\_1 deposited under accession number ATCC 207187;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;
- (h) a nucleotide sequence encoding a protein comprising the amino  
30 acid sequence of SEQ ID NO:152;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152, the fragment comprising eight contiguous amino acids of SEQ ID NO:152;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:151.

161. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:152;
- (b) a fragment of the amino acid sequence of SEQ ID NO:152, the fragment comprising eight contiguous amino acids of SEQ ID NO:152; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pe246\_1 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins.

162. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:153;
- (b) the nucleotide sequence of SEQ ID NO:153 from nucleotide 94 to nucleotide 1281;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone qf122\_3 deposited under accession number ATCC 207187;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qf122\_3 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:154;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:154, the fragment comprising eight contiguous amino acids of SEQ ID NO:154;

- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- 5 (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:153.

163. A protein comprising an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:154;
- (b) a fragment of the amino acid sequence of SEQ ID NO:154, the fragment comprising eight contiguous amino acids of SEQ ID NO:154; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
15 qf122\_3 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins.

164. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 20 (a) the nucleotide sequence of SEQ ID NO:155;
- (b) the nucleotide sequence of SEQ ID NO:155 from nucleotide 110 to nucleotide 742;
- (c) the nucleotide sequence of SEQ ID NO:155 from nucleotide 170 to nucleotide 742;
- 25 (d) the nucleotide sequence of the full-length protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;
- 30 (f) the nucleotide sequence of a mature protein coding sequence of clone qv538\_1 deposited under accession number ATCC 207187;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;

- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:156;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156, the fragment comprising eight contiguous amino acids of SEQ ID NO:156;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:155.

165. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:156;
- (b) a fragment of the amino acid sequence of SEQ ID NO:156, the fragment comprising eight contiguous amino acids of SEQ ID NO:156; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qv538\_1 deposited under accession number ATCC 207187;
- the protein being substantially free from other mammalian proteins.

166. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:157;
- (b) the nucleotide sequence of SEQ ID NO:157 from nucleotide 41 to nucleotide 757;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ys20\_1 deposited under accession number ATCC 207187;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ys20\_1 deposited under accession number ATCC 207187;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:158;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:158, the fragment comprising eight contiguous amino acids of SEQ ID NO:158;

5 (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and  
10 that has a length that is at least 25% of the length of SEQ ID NO:157.

167. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:158;  
15 (b) a fragment of the amino acid sequence of SEQ ID NO:158, the fragment comprising eight contiguous amino acids of SEQ ID NO:158; and  
(c) the amino acid sequence encoded by the cDNA insert of clone  
ys20\_1 deposited under accession number ATCC 207187;  
the protein being substantially free from other mammalian proteins.

20

168. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:159;  
(b) the nucleotide sequence of SEQ ID NO:159 from nucleotide 28 to  
25 nucleotide 2253;  
(c) the nucleotide sequence of SEQ ID NO:159 from nucleotide 568 to  
nucleotide 2253;  
(d) the nucleotide sequence of the full-length protein coding sequence  
of clone as180\_1 deposited under accession number ATCC XXXXXX;  
30 (e) a nucleotide sequence encoding the full-length protein encoded by  
the cDNA insert of clone as180\_1 deposited under accession number ATCC  
XXXXXX;  
(f) the nucleotide sequence of a mature protein coding sequence of  
clone as180\_1 deposited under accession number ATCC XXXXXX;

- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:160;
- 5 (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:160, the fragment comprising eight contiguous amino acids of SEQ ID NO:160;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees
- 10 C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:159.
- 15
169. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:160;
- (b) a fragment of the amino acid sequence of SEQ ID NO:160, the
- 20 fragment comprising eight contiguous amino acids of SEQ ID NO:160; and
- (c) the amino acid sequence encoded by the cDNA insert of clone as180\_1 deposited under accession number ATCC XXXXXX;
- the protein being substantially free from other mammalian proteins.

Fig. 1A




Fig. 1B







Fig. 2



## SEQUENCE LISTING

<110> Jacobs, Kenneth  
 McCoy, John M.  
 LaVallie, Edward R.  
 Collins-Racie, Lisa A.  
 Evans, Cheryl  
 Merberg, David  
 Treacy, Maurice  
 Agostino, Michael J.  
 Steininger II, Robert J.  
 Spaulding, Vikki  
 Wong, Gordon G.  
 Clark, Hilary  
 Fechtel, Kim  
 Genetics Institute, Inc.

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6075-83A

<140>

<141>

<160> 240

<170> PatentIn Ver. 2.0

<210> 1

<211> 1925

<212> DNA

<213> Homo sapiens

<400> 1

```

aggtcgtcac agacgatgat ggccaggccc cggaggctaa ggacggcagc tccttttagcg 60
gcagagtttt ccgagtgacc ttcttgatgc tggctgtttc tctcaccgtt cccctgcttg 120
gagccatgat gctgctggaa tctcctatag atccacagcc tctcagcttc aaagaacccc 180
cgctcttgct tgggtgttctg catccaaata cgaagctgcy acaggcagaa aggctgtttg 240
aaaatcaact tgttgaccg gagtccatag cacatattgg ggatgtgatg tttactggga 300
cagcagatgg cccggtcgta aaacttgaaa atggtgaaat agagaccatt gcccggtttg 360
gttcggggccc ttgcaaaacc cgagatgatg agcctgtgtg tgggagaccc ctgggtatcc 420
gtgcagggcc caatgggact ctctttgtgg ccgatgcata caagggacta tttgaagtaa 480
atccctggaa acgtgaagtg aaactgctgc tgcctccga gacaccatt gaggggaaga 540
acatgtcctt tgtgaatgat cttacagtca ctcaggatgg gaggaagatt tatttcaccg 600
attctagcag caaatggcaa agacgagact acctgttctt ggtgatggag ggcacagatg 660
acgggcgccct gctggagtat gatactgtga ccaggggaagt aaaagtttta ttggaccagc 720
tgcggttccc gaatggagtc cagctgtctc ctgcagaaga ctttgtcctg gtggcagaaa 780
caaccatggc caggatacga agctcttttag tcaagagacg gtgatgaagt ttgtgcccg 840
gtacagcctc gtcctagaac tcagcgacag cgggtgccttc cggagaagcc tgcagatcc 900
cgatgggctg gtggccacct acatcaccga ggtgcacgaa cagatgggc acctgtacct 960
gggctctttc aggtccccct tcctctgcag actcagcctc caggctgttt agccctccca 1020
gatagctgcc cctgccacgc aggccaggag tcttcacact caggcaccag gcctgggtcca 1080
ggaggagctg tgacacagt cgtggttcaa gtgtccacat gcacctgtta gtccctgaga 1140
gggtgtggga atggctgctt cattcctcga ggtgcccgg gccccacctg ggcttgtctt 1200
tctgtttaga ggaagtgtg acatatctgc catgaggaac ataaattcat gtaaaagccat 1260
tttctcttaa aaaaaacaa actttctaa tacaatcatt ctctaggatt tgggaagctc 1320
cttgcaactg gaacagggtc cagggtgggtg gagcagtaag gcactacca gagagcttgc 1380
tgctgcggcc ctgtcctgcg gcctcaagt tcttctttac tatatataac gtgcggtcat 1440
acctttcttc gttgtggtgg ggatggaaga gcagaggag catggcccag ggggtttgag 1500
gccagcggtg agagccgtgt tagccaagac atggaactgt gttctcaagg gttatgtggg 1560

```